

Performances, meat quality and boar taint of castrates and entire male pigs fed a standard and a raw potato starch-enriched diet

C. Pauly^{1,2}, P. Spring¹, J. V. O'Doherty², S. Ampuero Kragten³ and G. Bee^{3†}

¹Department of Animal Science, Swiss College of Agriculture, Zollikofen 3052, Switzerland; ²School of Agriculture, Food Science and Veterinary Medicine, University College Dublin, Dublin 4, Ireland; ³Agroscope Liebefeld Posieux Research Station ALP, Posieux 1725, Switzerland

(Received 30 March 2007; Accepted 9 June 2008; First published online 22 July 2008)

In Europe there is increasing concern about the common practice of surgical castration of piglets without anaesthesia. One possible alternative to completely avoid castration is entire male pig production. Thus, the objective of the study was to compare the growth performance, carcass characteristics, organ weights, meat quality traits, fat score and boar taint compounds in the adipose tissue of group-penned entire male pigs and castrates. Furthermore, the effect of raw potato starch (RPS) fed for 7 days prior to slaughter was determined. Pigs (n = 36) were blocked by BW into 12 blocks (3 littermates/block) and assigned to three experimental groups: surgical castrates (C); entire males (EM); and entire males offered RPS (30 g RPS/100 g diet) for 7 days prior to slaughter (EM+). Pigs had ad libitum access to the feed from 22 to 107 kg, individual feed intake was recorded daily and BW once a week. Entire males grew slower (EM: 771, EM+: 776 v. C: 830 g/day; $P < 0.01$), consumed less feed (EM: 1.87, EM+: 1.89 v. C: 2.23 kg/day; $P < 0.01$) and were more efficient (feed conversion ratio: EM: 2.42, EM+: 2.44 v. C: 2.69 kg/kg; $P < 0.01$) than C. Compared to C, carcass dressing percentage was lower (EM: 79.4, EM+: 79.4 v. C: 81.6%; $P < 0.01$) and percentage of valuable cuts was higher (EM: 57.3, EM+: 56.5 v. 52.6%; $P < 0.01$) in entire males. The hearts (EM: 426, EM+: 425 v. C: 378 g), kidneys (EM: 387, EM+: 378 v. C: 311 g), bulbourethral (EM: 200, EM+: 195 v. C: 7 g) and salivary glands (EM: 99, EM+: 94 v. C: 42 g) were heavier ($P < 0.001$) in entire males than in C. Meat quality traits did not ($P > 0.05$) differ among experimental groups but the adipose tissue was more unsaturated in entire males than in C as indicated by the higher fat scores (EM: 69.1, EM+: 67.2 v. C: 63.6; $P < 0.01$). Feeding RPS reduced ($P = 0.04$) the skatole tissue concentrations (expressed in $\mu\text{g/g}$ lipid) in EM+ (0.22) compared to EM (0.85), whereas androstenone and indole levels were not ($P \geq 0.60$) affected (EM: 1.7 and 0.10, EM+: 2.0 and 0.09, respectively). Although the current results confirmed the high efficiency of entire males compared to castrates, the observed high androstenone levels represent a major challenge to implement entire males production.

Keywords: carcass characteristics, entire male pig, growth performance, raw potato starch, skatole

Introduction

The Swiss Parliament has passed a law prohibiting castration of piglets without anaesthesia taking effect in 2009. This new situation challenges the production chain in finding the best-adapted alternative for the Swiss market. Fattening entire male pigs could be one solution. However, boar taint, an unpleasant odour and flavour can develop in the meat from some entire male pigs. Androstenone (5 α -androst-16-en-3-one) and skatole (3-methyl-indole) are thought to be the key contributors to boar taint (Bonneau

et al., 2000). Rearing entire males instead of castrates has a number of advantages including higher efficiency, leaner carcasses and lower faecal and urinary nitrogen losses (Bonneau, 1998). It also offers welfare advantages due to the absence of castration pain and stress (Prunier *et al.*, 2006). Results of some studies also report higher growth rates for entire males than for C (Andersson *et al.*, 1997; Turkstra *et al.*, 2002). However, responses of anabolic steroids production by entire males are inconsistent because performance differences between entire males and castrates depend on genotype and housing conditions. The available results on growth performance are based on older genotypes and also on individually penned animals and,

[†] E-mail: Giuseppe.Bee@alp.admin.ch

therefore, might not fully reflect the present situation under commercial conditions (Dunshea *et al.*, 2001; Suster *et al.*, 2006). Compared to castrates (Cronin *et al.*, 2003) and gilts (Rydhmer *et al.*, 2006), entire males are more aggressive and sexually active and their fighting and mounting behaviour can result in leg problems and skin damage.

Androstenone is synthesized in the testes of sexually mature boars and released into the blood, while skatole is formed by intestinal bacteria from L-tryptophan and absorbed into the blood (Claus *et al.*, 1994). Both substances accumulate in the adipose tissue. Thus, their tissue concentrations are related to sexual maturity at the time of slaughter (Babol *et al.*, 2004; Zamaratskaia *et al.*, 2004). Feeding fermentable carbohydrates such as raw potato starch (RPS) has been shown to strongly reduce skatole concentration in the adipose tissue of barrows (Claus *et al.*, 2003) and entire male pigs (Andersson *et al.*, 2005; Zamaratskaia *et al.*, 2005a). As skatole breakdown in the adipose tissue occurs rapidly, when microbial formation is reduced, feeding resistant starch during the week prior to slaughter seems sufficient to achieve significant reductions (Lösel and Claus, 2005). These authors suggested a dietary incorporation of 30% RPS to be a good compromise between economic considerations and the achieved reduction in the skatole concentration in the adipose tissue.

In Switzerland, the last trials aiming to evaluate performances and meat quality of entire males were conducted more than 20 years ago (Stoll, 1982; Schnegg *et al.*, 1985). Since then, genetics improvements occurred, in particular carcass leanness increased markedly. As the carcass composition has a major impact on adipose tissue composition (Bee *et al.*, 2002), which is part of the routine carcass quality evaluation in Switzerland and is also relevant for the market price of the carcass, changing the production system to fattening entire males would likely require dietary adaptations to correct for those changes. There is also a lack of scientific investigations regarding androstenone and skatole levels in the adipose tissue from Swiss pigs.

Thus, the aims of the present study were to evaluate performance, carcass composition, meat quality and boar taint of entire male pigs compared to castrates reared under Swiss environment- and welfare-friendly housing conditions. In addition, the effect of RPS feeding for 1 week prior to slaughter was evaluated.

Material and methods

Growth trial and carcass measurements

The Swiss Federal Committee for Animal Care and Use approved all procedures involving animals.

The experiment was conducted from February to July 2006 and included a total of 36 Swiss Large White male pigs originating from 12 litters. From weaning until the start of the experimental period (mean \pm s.e. BW = 21.6 \pm 0.3 kg; age = 65 \pm 0.9 days), pigs were group-penned and had *ad libitum* access to a commercial starter diet. At an average BW of 13.1 \pm 0.4 kg, the littermates were allocated to

Table 1 Composition of the growing and finishing diet (as-fed basis)

	Grower diet	Finisher diet	
		Without RPS	With RPS
Barley	55.1	24.8	26.0
Wheat	24.1		10.0
Wheat starch		30.0	
RPS			30.0
Sugar beet pulp		10.0	10.0
Wheat bran	3.4	8.0	6.0
Potato protein	7.1		4.4
Rape cake	3.9	0.7	
Linseed cake		8.2	
Soybean cake		7.3	2.5
Corn gluten		5.6	6.5
Sugar beet molasses	3.0	3.0	
Mix fat 50/50			1.78
Lysine-HCl	0.33	0.312	0.276
DL-Methionine	0.038	0.008	0.004
L-Threonine	0.066	0.034	0.002
L-Tryptophane	0.008		0.022
Dicalcium phosphate	0.692	0.404	0.866
Sodium chloride	0.44	0.046	0.256
Pellan ¹	0.3	0.3	0.3
Calcium carbonate	1.126	0.896	0.694
Vitamin-mineral premix ²	0.4	0.4	0.4
Analysed composition (in the DM)			
CP (%)	18.7	16.7	16.8
Lysine (%)	11.7	9.1	9.5
Lysine/DE (g/MJ)	0.8	0.6	0.6
Crude fat (%)	2.0	2.6	3.8
Crude fibre (%)	4.1	5.2	4.3
Ca (%)	0.82	0.69	0.72
P (%)	0.63	0.53	0.56
Calculated energy content ³			
DE (MJ/kg DM)	14.9	14.9	15.6

RPS = raw potato starch; DE = digestible energy; DM = dry matter.

¹Pellan = a binder that aids in pellet formation (Mikro-Technik GmbH & Co., KG, Germany).

²Vitamin-mineral premix supplied the following nutrients per kg of diet: 20 000 IU vitamin A, 200 IU vitamin D₃, 39 IU vitamin E, 2.9 mg riboflavin, 2.4 mg vitamin B₆, 0.010 mg vitamin B₁₂, 0.2 mg vitamin K₃, 10 mg pantothenic acid, 1.4 mg niacin, 0.48 mg folic acid, 199 g cholin, 0.052 mg biotin, 52 mg Fe as Fe-sulphate, 0.16 mg I as Ca(IO₃), 0.15 mg Se as Na₂Se, 5.5 mg Cu as CuSO₄, 81 mg Zn as ZnO₂, 15 mg Mn as MnO₂.

³DE content calculated according to the formula: DE = 18.974 \times CP (g/g DM) + 33.472 \times crude fat (g/g DM) + 16.611 \times NFE (g/g DM) - 21.216 \times crude fibre (g/g DM) + 16.611 \times NFE (g/g DM). NFE = DM - ash - CP - crude fat - crude fibre.

the three experimental groups according to BW. One group of 12 animals was surgically castrated (C). From 21.6 to 65 kg BW, and from 65 to 105.8 kg BW, the entire males and C had *ad libitum* access to the same grower and finisher diet (Table 1). For the last 7 days prior to slaughter, one group of entire males (EM+; n = 12) received a finisher diet with 30% RPS (Emsland-Stärke GmbH, D-49824 Emlichheim, Germany), whereas C and the remaining entire males (EM; n = 12) were kept on the finisher diet without RPS. The RPS was replaced with wheat starch and special

care was taken to maintain a similar nutrient composition. All diets were pelleted (4.5-mm diameter) at 60°C.

The three group-pens in which C, EM and EM+ were reared were equipped with single-space computerized feeders as previously described (Bee *et al.*, 2008). Pigs were weighed weekly and feed intake was recorded daily. From 80 kg BW until slaughter (BW = 105.8 ± 0.9 kg), pens were cleaned daily and barn ventilation set at maximum power to reduce skatole absorption through the skin and the lungs (Hansen *et al.*, 1994). Feed was withheld 15 h from the pigs before transportation to the research station abattoir where they were electrically stunned, exsanguinated, scalded, mechanically de-haired and eviscerated. Internal organs were removed and hot carcass weight was obtained. The weight of the testes, bulbourethral and salivary glands as well as the heart, liver and kidneys were assessed. Furthermore, a fat sample over the croup was taken for the determination of the fat score according to the method previously described (Scheeder *et al.*, 1999). Thirty minutes after exsanguination, the carcasses entered the air-chilling system (3°C) for 24 h. One day after slaughter, the left side of each carcass was weighed and dissected according to meat cutting standards applied by the Swiss Performance Testing Station (MLP, Sempach, Switzerland), as previously described by Bee (2001).

Objective meat quality measurements

The pH of the *longissimus* muscle (LM) at 30 min and 24 h *post mortem* was measured using a pH meter (pH196-S, WTW, Weilheim, Germany) equipped with an electrode (Eb4; WTW) and a temperature probe. Sets of measurements were obtained at the 10th rib of the right carcass side at 30 min by insertion of the pH and temperature probe between the ribs from the inside of the carcass.

At 24 h *post mortem*, the LM from the 10th to 13th rib level was excised and pH was determined at the 10th rib level. Subsequently, four 1.5-cm-thick LM chops were cut and labelled A, B, C and D. From chops A and C, drip loss was determined as the amount of purge formed during storage of chops at 4°C for 48 h (Honikel, 1998). The A and C samples were then vacuum-packed and stored at -20°C. From chops B and D, light reflectance coordinates (L^* : lightness, a^* : redness and b^* : yellowness) of the muscle surface were determined following a 10 min bloom, using a Chroma Meter CR-300 with a D65 light source (Minolta, Dietikon, Switzerland). After colour measurements were made, the samples were vacuum-packed and stored at -20°C until Warner-Bratzler shear force was determined.

Frozen samples were thawed for 24 h at 2°C, subsequently kept at room temperature for 1 h and the thaw loss was determined. Both samples were then cooked on a grill plate (Beer Grill AG, Zurich, Switzerland) at 190°C to 195°C to an internal temperature of 69°C, and cooking losses were measured. Using a Warner-Bratzler shear device (Model 3000; G-R Electric Mfg. Co., Manhattan, KS, USA), shear force values were determined as previously described (Bee *et al.*, 2007). In addition, adipose tissue (both layers)

was sampled at the 10th to 13th rib level, vacuum-packed and stored at -20°C until further analysis.

Analysis of skatole and androstenone in adipose tissue and in plasma

Androstenone, skatole and indole concentrations in the adipose tissue were analysed according to the method described by Hansen-Moller (1994). Briefly, adipose tissue samples were liquefied in a microwave oven for 2×2 min at 250 W. The liquefied lipids were centrifuged for 2 min at $11\,000 \times g$ at 20°C and kept at around 47°C. The water was then removed and 0.50 ± 0.01 g water-free liquid fat was placed in 2.5 ml eppendorf tubes in duplicates and an internal standard was added (1 ml methanol containing 0.496 mg/l androstanone and 0.050 mg/l 2-methylindole). After vortexing for 30 s, the tubes were incubated for 5 min at 30°C in an ultrasonic-water bath (RK 255 H; Bandelin Sonorex, Zürich, Switzerland), kept at 0°C on ice for 20 min and then centrifuged for 20 min at $11\,000 \times g$ at 4°C. Finally, 50 µl of the supernatant was transferred into a high-performance liquid chromatography (HPLC) vial for androstenone, skatole and indole analysis with an HPLC system from Hewlett-Packard series 1100. The procedure of the analysis with the HPLC is similar to the one described by Hansen-Moller (1994). Concentrations were expressed per g of the lipid fraction from adipose tissue. The detection limits were 0.3 µg for androstenone and 0.03 µg for skatole and indole. Blood samples from pigs were collected at exsanguination, centrifuged at $840 \times g$ for 15 min and plasma stored at -20°C until analysis. Subsequently, skatole and indole levels in the plasma were determined by HPLC after ether extraction using the method described by Lösel (2006).

Statistical analysis

Data for growth performance, carcass characteristics and meat quality traits were analysed with the MIXED procedure of SAS (v. 9.1; SAS Institute, Cary, NC, USA). The model used included litter and experimental groups as fixed effects. The individual pig was the experimental unit for analysis of all data. Least square means were separated using the PDIF option ($P < 0.05$). Tendencies were established at a $P < 0.10$. Androstenone, skatole and indole concentrations in the adipose tissue and plasma were analysed with the non-parametric Mann-Whitney *U* test with the statistical software package NCSS (version 2007; NCSS, Kaysville, Utah, USA). Two-tail test for androstenone and indole and one-tail test for skatole, right-tail in favour of EM+, were considered. Spearman's Rank correlation coefficients were determined between androstenone, skatole and indole levels in the adipose tissue and plasma and the organ weights.

Results

Growth, carcass characteristics and organ weights

Despite the late time point of castration, growth from weaning until the start of the experiment did not ($P = 0.37$)

Table 2 Growth performance, carcass characteristics and organ weights from castrates (C) and entire males fed a finisher diet without (EM) or with raw potato starch in the diet (EM+)

	C	EM	EM+	s.e.	P-values
Initial BW (kg)	21.9	21.9	21.2	0.3	0.18
BW at slaughter (kg)	105.7	106.0	105.7	0.9	0.95
ADFI (kg/day)					
Grower period	1.93 ^a	1.67 ^b	1.65 ^b	0.05	0.001
Finisher period	2.56 ^a	2.07 ^b	2.15 ^b	0.05	<0.001
Grower–finisher period	2.23 ^a	1.87 ^b	1.89 ^b	0.04	<0.001
ADG (g/day)					
After weaning period	379	360	363	10	0.37
Grower period	828 ^a	767 ^b	765 ^b	16	0.01
Finisher period	831	776	791	22	0.21
Grower–finisher period	830 ^a	771 ^b	776 ^b	14	0.01
Feed conversion ratio (kg/kg)					
Grower period	2.33 ^a	2.17 ^b	2.15 ^b	0.03	0.001
Finisher period	3.07 ^a	2.69 ^b	2.73 ^b	0.05	<0.001
Grower–finisher period	2.69 ^a	2.42 ^b	2.44 ^b	0.03	<0.001
Age at slaughter (days)	167.7 ^a	176.3 ^b	175.3 ^b	2.3	0.03
Carcass characteristics					
Hot carcass weight (kg)	86.3 ^d	84.1 ^{de}	83.9 ^e	0.8	0.08
Carcass yield (%) ¹	81.6 ^a	79.4 ^b	79.4 ^b	0.3	<0.001
Lean meat (%) ²	52.6 ^a	57.3 ^b	56.5 ^b	0.4	<0.001
Loin (%)	23.8 ^a	25.4 ^b	25.3 ^b	0.2	<0.001
Ham (%)	17.2 ^a	18.9 ^c	18.3 ^b	0.2	<0.001
Shoulder (%)	11.6 ^a	13.0 ^b	12.9 ^b	0.1	<0.001
Belly (%)	17.8 ^a	16.9 ^b	17.4 ^{ab}	0.2	0.04
Subcutaneous fat (%) ³	15.6 ^a	12.0 ^b	12.4 ^b	0.3	<0.001
Omental fat (%) ⁴	2.3 ^a	1.2 ^b	1.3 ^b	0.1	<0.001
Backfat thickness (mm)	22.1 ^a	13.0 ^b	13.3 ^b	1.0	<0.001
Organ characteristics					
Heart (g)	378 ^a	426 ^b	425 ^b	8	<0.001
Liver (g)	1630	1743	1672	48	0.27
Kidney (g)	311 ^a	387 ^b	378 ^b	11	<0.001
Testes (g)		756	712	28	0.28
Bulbourethral gland (g)	7 ^a	200 ^b	195 ^b	14	<0.001
Bulbourethral gland length (cm)	5.3 ^a	13.3 ^b	13.4 ^b	0.4	<0.001
Salivary gland (g)	42 ^a	99 ^b	94 ^b	7	<0.001

ADFI = average daily feed intake; ADG = average daily gain.

^{a–c}Significant effect of the experimental groups ($P < 0.05$).^{d,e}Significant effect of the experimental groups ($P < 0.10$).^P-values = probability values for the experimental groups.¹Carcass yield = hot carcass weight as percentage of the BW at slaughter.²Lean meat percentage = sum of denuded shoulder, loin and ham weights as percentage of cold carcass weight.³Subcutaneous fat = sum of external fat from the shoulder, loin and ham expressed as percentage of cold carcass weight.⁴Omental fat percentage = weight of the omental fat as percentage of cold carcass weight.

differ among the experimental groups (Table 2). From 21.6 kg BW until slaughter, EM and EM+ grew slower ($P < 0.01$), consumed less feed ($P < 0.001$) and were more efficient ($P < 0.001$) than C. The differences for average daily feed intake (ADFI), average daily gain (ADG) and feed conversion ratio (FCR) were already apparent during the grower period. Because of the slower growth rate, EM and EM+ were 8.1 days older ($P = 0.03$) at slaughter than C. Feeding RPS the last week before slaughter did not affect overall growth performance (data not shown).

Carcass yield was lower ($P < 0.001$) and lean meat percentage was markedly higher ($P < 0.001$) in EM and EM+ than C (Table 2). In accordance, subcutaneous fat

percentage and backfat thickness were lower ($P < 0.001$) in carcasses of EM and EM+ compared to C. The greater carcass leanness resulted from the higher ($P < 0.001$) loin, ham and shoulder percentages in entire male carcasses. Compared to C, the carcasses of EM had lower ($P = 0.04$) belly percentage with intermediate values for EM+. Entire males had heavier hearts and kidneys compared to castrates ($P < 0.001$ for both; Table 2). Bulbourethral and salivary glands were also heavier and bulbourethral gland was longer in EM and EM+ compared to C ($P < 0.001$ for each). The dietary RPS did not ($P > 0.10$) affect the weights of the evaluated organs as no differences were observed between EM and EM+.

Meat and backfat quality

The effects of the experimental groups on quality traits of the LM were minor (Table 3). The muscle temperature 30 min *post mortem* of the EM and EM+ was lower ($P = 0.02$) than that of C. The LM from EM and EM+ had numerically higher ($P = 0.16$) a^* values and lost more ($P = 0.14$) purge during thawing than the LM of C. Feeding RPS the last week before slaughter did not ($P > 0.10$) affect objective meat quality traits as no differences were observed between EM and EM+. The fat score was markedly higher ($P < 0.001$) in the adipose tissue of EM and EM+ compared to C, indicating a higher degree of unsaturation. The fat score was closely related to carcass fatness as suggested by the negative correlations between fat score and subcutaneous fat percentage ($r = -0.81$, $P < 0.001$) and croup fat thickness ($r = -0.70$, $P < 0.001$).

Table 3 Meat quality traits determined in the longissimus muscle and fat score analysed in the backfat from castrates (C), and entire males fed a finisher diet without (EM) or with raw potato starch in the diet (EM+)

	C	EM	EM+	s.e.	P-values
Initial pH	6.16	6.15	6.14	0.05	0.91
Initial temperature (°C)	39.2 ^a	39.8 ^b	39.8 ^b	0.2	0.02
Ultimate pH	5.60	5.57	5.56	0.02	0.38
L^*	50.5	50.0	50.6	0.4	0.65
a^*	6.2	6.7	6.8	0.2	0.16
b^*	2.8	3.0	3.2	0.2	0.52
Drip loss (%)	4.0	4.7	4.9	0.4	0.34
Thawing loss (%)	8.3	9.5	9.4	0.5	0.14
Cooking loss (%)	13.1	13.9	14.4	0.6	0.38
Total loss (%)	20.3	22.1	22.4	0.8	0.13
Shear force (kg)	4.99	5.34	5.46	0.18	0.19
Loin purge (%)	0.58	0.59	0.58	0.06	0.98
Neck purge (%)	0.60	0.77	0.52	0.11	0.32
Fat score	63.6 ^a	69.1 ^b	67.2 ^b	0.6	<0.001

^{a,b}Significant effect of the experimental groups ($P < 0.05$).
P-values = probability values for the experimental groups.

Androstenone and skatole concentrations

Androstenone concentrations in the adipose tissue of all C were below the detectable limits of the used method (Table 4). The androstenone concentrations did not ($P = 0.73$) differ in the adipose tissue of EM and EM+. A high variability in the androstenone concentration was observed ranging from below detection limits up to 4 and 6.1 µg/g of the lipid fraction from adipose tissue in EM and EM+, respectively. However, skatole concentrations were lower (−74%; $P = 0.044$) in EM+ than in EM, indicating a marked effect of dietary RPS supply. The lowest skatole concentrations were observed in C. Nevertheless, it is worthwhile mentioning that in the adipose tissue of one C skatole concentration was, with a concentration of 0.24 µg, high. In accordance with the findings in the adipose tissue, the average skatole concentration in the plasma from EM+ tended to be lower ($P = 0.09$) than for EM. The skatole concentrations in the adipose tissue and plasma were highly positively correlated ($r = 0.91$, $P < 0.001$). Although the indole concentrations in both the adipose tissue and plasma were not ($P \geq 0.26$) affected by the diet, the concentrations of this compound were highly correlated ($r = 0.81$, $P < 0.001$) between adipose tissue and plasma. In contrast, androstenone and skatole concentrations in the adipose tissue were not ($r = 0.24$, $P = 0.46$) related.

The androstenone concentration in the adipose tissue was negatively correlated with the percentage subcutaneous fat ($r = -0.57$, $P < 0.001$) and backfat thickness ($r = -0.51$, $P = 0.001$). Similarly, skatole concentrations were also correlated with both traits (percentage subcutaneous fat: $r = -0.48$, $P < 0.01$ and backfat thickness: $r = -0.46$, $P < 0.01$). Furthermore, with increasing androstenone concentrations in the adipose tissue, the weight of the salivary ($r = 0.63$, $P \leq 0.001$) and bulbourethral gland ($r = 0.67$, $P \leq 0.001$) increased as well as the bulbourethral gland was longer ($r = 0.49$, $P = 0.02$). In contrast, no correlations were found between androstenone concentrations in the adipose tissue and the testes weight

Table 4 Skatole and indole levels in the adipose tissue and in plasma and androstenone concentrations in the adipose tissue from castrates (C) and entire males fed a finisher diet without (EM) or with raw potato starch in the diet (EM+)

		C	EM	EM +	P-values
Adipose tissue [†]					
Androstenone (μg)	Mean	≤0.3	1.7	2.0	0.73
	Min.–max.		≤0.3–4.0	0.5–6.1	
Skatole (μg)	Mean	0.07	0.85 ^a	0.22 ^b	0.04
	Min.–max.	0.03–0.24	0.06–3.34	≤0.03–0.64	
Indole level (μg)	Mean	0.03	0.10	0.09	0.60
	Min.–max.	≤0.03–0.04	≤0.03–0.51	≤0.03–0.25	
Plasma					
Skatole level (ng/ml)	Mean	1.7	20.0	7.0	0.09
	Min.–max.	0.6–10.2	0.8–89.9	0.2–29.8	
Indole level (ng/ml)	Mean	1.0	6.0	8.1	0.26
	Min.–max.	0.6–1.6	0.6–36.9	0.6–29.1	

^{a,b}Significant effect of the experimental groups EM and EM+ ($P < 0.05$).

[†]Androstenone, skatole and indole concentrations in the adipose tissue are expressed as µg/g liquefied lipids.

($r = 0.35$, $P = 0.09$). The correlation coefficients between skatole or indole concentrations and the aforementioned organ weights and length were not significant ($P > 0.05$).

Discussion

Differences in performance, carcass characteristics, meat quality and boar taint level between entire males and castrates were reported to depend on a variety of factors including genetics, nutrition and housing conditions (Barton-Gade, 1987; Bonneau, 1998). However, in particular, the reported differences in feed intake and weight gain have been inconsistent (Xue *et al.*, 1997). In this study, EM and EM+ consumed markedly less feed than C. First, this could be due to a negative influence of sexual hormones on appetite. Claus and Weiler (1987) reported that oestrogen and androgen influenced negatively feed intake. They showed that in castrates, the application of an implant of testosterone reduced feed intake by about 25%. Second, the lower ADFI could be related to the observed higher physical activity of EM and EM+, which was not limited to the finisher period, but both EM and EM+ already showed more mounting and aggression at the end of the grower period. In agreement, several authors (Giersing *et al.*, 2000; Cronin *et al.*, 2003; Rydhmer *et al.*, 2006) reported that entire males spent less time eating and more time mounting, a behaviour observed primarily after the onset of sexual maturity. Suster *et al.* (2006) suggested that such changes in behaviour could also induce social stress, which could stimulate cortisol production. Elevated cortisol levels have been shown to reduce feed intake, decrease production of growth hormone and IGF-1 (Black *et al.*, 2001). The observed consequences of the reduced feed intake were not surprising. In fact EM and EM+ gained on average less than C (ADG: -54 g/day), but were still more efficient (FCR: -0.26 kg/kg). These findings are in line with recent studies reporting a tendency towards lower ADG for entire males reared in group compared to castrates (Dunshea *et al.*, 2001; D'Souza and Mullan, 2002; Suster *et al.*, 2006). Contrarily, some previous studies reported that entire male pigs had higher ADG and better FCR than castrates (Whittemore *et al.*, 1988; Campbell *et al.*, 1989; Dunshea *et al.*, 1993). This might be surprising as the genetic growth potential was markedly improved in the last decade due to breeding efforts. However, compared to recent experiments, these studies with entire males of the nineties were carried out with individually penned pigs, which limited the social interactions and, thus, lowered the energy expenditure on activities unrelated to growth.

Carcass yield and carcass fat deposition were lower in EM and EM+ than in C, which confirms the results reported by Sather *et al.* (1999), Dunshea *et al.* (2001) and Zeng *et al.* (2002). It is well known that lower backfat deposition rate affects lipid composition of the adipose tissue as the degree of unsaturation of the lipids increases (Bee *et al.*, 2002). In accordance, EM and EM+ had higher fat scores than C and the fat score was negatively correlated with

backfat thickness. These findings are in line with earlier studies of Barton-Gade (1987) who determined higher iodine value in the backfat of entire males than castrates.

While general meat quality parameters were not affected by treatment, specific parameters involved in boar taint did markedly differ. Boar taint is mainly caused by the accumulation of androstenone and skatole or both in the adipose tissue. Androstenone is produced in the testes and its concentration increases with sexual maturity (Bonneau, 1982). Comprehensive studies have shown that between 18 and 42% of the entire males have androstenone levels above 1.0 $\mu\text{g/g}$ adipose tissue (Walstra *et al.*, 1999; Bonneau *et al.*, 2000). The androstenone concentration of 1 $\mu\text{g/g}$ adipose tissue has been proposed to be the threshold value for consumer acceptability in the EU (Desmoulin *et al.*, 1982; Bonneau *et al.*, 1992; Annor-Frempong *et al.*, 1997). When this value is expressed on the basis of the lipid fraction from the adipose tissue only, as requested by the analytical procedure used in this study, the concentration is 1.7 μg . This relation between both methods was determined after analysis of samples with both extraction methods (lipid fraction from the adipose tissue only and adipose tissue). The average androstenone concentrations reported in the present trial are just at or above the aforementioned threshold level. Very limited information is available on androstenone and skatole concentrations in the Swiss pig population. In a recent large field trial, Pauly *et al.* (2007) showed that only 3.8% of the analysed adipose tissue samples ($n = 170$) from entire males had androstenone levels above 1.7 $\mu\text{g/g}$ of the lipid fraction from adipose tissue. Fredriksen *et al.* (2006) reported that the herd average androstenone level varied considerably in three herds in Norway (expressed per g adipose tissue: means from 0.66 to 1.67 μg ; and expressed per g of the lipid fraction from the adipose tissue: 1.12 to 2.83 μg), even when adjusted for fattening system, season, breed and carcass weight. This suggests that feeding strategies, ADG, and also genetic and environment (e.g. photoperiod) are factors that can influence puberty and, thus, affect the development of boar taint. Sexual maturity of entire male pigs is difficult to determine because it is a continuously developing process (Malmgren, 1993). However, in accordance to Andersson *et al.* (1999), the weights of the testes and bulbourethral glands indicated that the majority of the EM and EM+ had attained puberty at slaughter. Furthermore, animals born earlier, which were heavier and slaughtered first, showed a tendency for higher androstenone levels in the adipose tissue. Giersing *et al.* (2000) reported that the level of plasma androstenone was related to rank order: high rank is often associated with higher BW. As heavier pigs were reaching sexual maturity earlier, they could have stimulated sexual development and androstenone secretion in other pen mates, which could explain the overall high androstenone concentrations in the EM and EM+ of this study. Besides the sexual organs, heart and kidney weights of EM and EM+ were also higher than for C. To our knowledge, no information has been reported

about such differences in pigs. In rats, blood pressure and the degree of cardiac hypertrophy were decreased by castration and increased by testosterone substitution (Morano *et al.*, 1990). Implanted steers with estradiol benzoate, progesterone and then with estradiol, trenbolone acetate, had higher circulating IGF-1 concentrations compared to bulls, and with it a significantly higher heart weight at slaughter (Schoonmaker *et al.*, 2002).

Skatole is a bacterial metabolite of tryptophan synthesized in the large intestine. The production of skatole greatly depends on the composition of the diet and the intestinal microflora (Jensen *et al.*, 1995; Claus *et al.*, 2003). Important quantities of butyrate are produced when RPS is fermented in the large intestine (Martin *et al.*, 2000; Mentschel and Claus, 2003), which reduces epithelial cell apoptosis (Claus *et al.*, 2003). Because tryptophan from cell debris is a substrate for skatole formation, lower skatole tissue and plasma levels can be expected. In accordance, a marked decrease in skatole content in the adipose tissue, and a tendency towards lower plasma levels were observed in EM+ compared to EM. Nevertheless, three animals on the RPS diet had skatole levels in the adipose tissue above the sensory threshold of 0.25 µg/g lipid fraction of the adipose tissue (0.16 µg/g adipose tissue), suggesting marked differences in the individual response to the diet. Genetic and dietary differences are thought to be two of the main factors explaining the sex-dependent and individual variation in the carcass adipose tissue skatole concentration of pigs. Skatole levels increase with puberty (Babol *et al.*, 2004; Zamaratskaia *et al.*, 2005b), so that sexual maturity may affect the skatole levels (Babol *et al.*, 1998b). Furthermore, there are important indications that genetic factors influence the skatole level in pigs. These include differences in skatole level between breeds (Babol *et al.*, 2004), intermediate heritability estimates for skatole in the adipose tissue (Bergsma *et al.*, 2007) as well as genetic differences in activities of cytochrome P450 (CYP), isoforms 2E1 and 2A6, which are two of the key enzymes responsible for skatole breakdown in the liver (Babol *et al.*, 1998a; Diaz and Squires, 2000; Lin *et al.*, 2006).

In this study, RPS was distributed only one week before slaughter, because RPS is an expensive nutritive compound. In contrast, other researchers offered RPS for 2 or 3 weeks (Andersson *et al.*, 2005; Lösel and Claus, 2005; Zamaratskaia *et al.*, 2005a) and obtained a complete wash-out of skatole from the adipose tissue; however, skatole concentration in the adipose tissue of control entire males was lower than in our study. Claus *et al.* (2003) measured a marked reduction in the adipose tissue accompanied by minimal skatole levels in the faeces and very low levels in the plasma by day four after RPS feeding. The skatole breakdown in the liver can be efficient and fast as the half-life of skatole in the adipose tissue is about 12 h (Hansen *et al.*, 1994). Thus, feeding RPS only one week before slaughter should be sufficient to decrease skatole concentration in the fat to the genetic level. As feeding RPS decreased skatole in adipose tissue, a reduction in indole could also be expected. In the present study,

no effects of RPS on indole concentrations in the adipose tissue and plasma were observed. Indole synthesis in the large intestine is mediated by many types of bacteria, whereas skatole production requires the presence of highly specific bacterial populations. Feeding RPS affected skatole-producing bacteria; however, indole-producing bacteria remain unaffected (Chen *et al.*, 2007).

In conclusion, the present results confirm that RPS supply was efficient in reducing skatole tissue concentrations. However, the androstenone concentrations in the adipose tissue of Swiss Large White entire males reared in a production system with high welfare standards were on average higher than the suggested threshold values for consumer acceptability in the EU. Meat quality did not differ among sexes but the adipose tissue composition might cause technological problems for further meat processing due to the high degree of unsaturation. Thus, specific feeding strategies need to be studied for entire male production. Besides the high androstenone concentrations, the lower ADG and in particular the low ADFI will pose a major challenge to implement fattening entire males instead of castrates. No conclusive result was obtained to show whether the lower ADFI was due to more active and aggressive behaviour or a general low ingestion capacity of entire male pigs.

Acknowledgements

The authors would like to thank C. Biolley, B. Dougoud, S. Dubois, G. Guex, G. Maïkoff and P. Stoll from Agroscope Liebefeld Posieux Research Station ALP (Posieux, Switzerland) for their support and technical assistance.

Prof. Dr R. Claus and Dr M. Lacorn from the University Hohenheim (Stuttgart, Germany) are gratefully acknowledged for technical support and determination of skatole levels in the plasma and UFAG-Laboratorien (Sursee, Switzerland) for determination of the fat score.

This trial was a part of the project 'ProSchwein'. Project ProSchwein is acknowledged for supporting the present study. We thank the Federal veterinary office, Migros, Suisseporcs, Coop, Suisag, Federal Office for Agriculture, Proviande and Swiss animal protection for financial support.

References

- Andersson K, Schaub A, Andersson K, Lundström K, Thomke S and Hansson I 1997. The effects of feeding system, lysine level and gilt contact on performance, skatole levels and economy of entire male pigs. *Livestock Production Science* 51, 131–140.
- Andersson HK, Hullberg A, Malmgren L, Lundström K, Rydhmer L and Squires J 1999. Sexual maturity in entire male pigs: environmental effects, relations to skatole level and female puberty. *Acta Agriculturae Scandinavica, Section A – Animal Sciences* 49, 103–112.
- Andersson HK, Andersson K, Zamaratskaia G, Rydhmer L, Chen G and Lundström K 2005. Effect of single-sex or mixed rearing and live weight on performance, technological meat quality and sexual maturity in entire male and female pigs fed raw potato starch. *Acta Agriculturae Scandinavica, Section A – Animal Sciences* 55, 80–90.
- Annor-Frempong IE, Nute GR, Whittington FW and Wood JD 1997. The problem of taint in pork: I. Detection thresholds and odour profiles of androstenone and skatole in a model system. *Meat Science* 46, 45–55.

- Babol J, Squires EJ and Lundström K 1998a. Hepatic metabolism of skatole in pigs by cytochrome P450E1. *Journal of Animal Science* 76, 822–828.
- Babol J, Squires EJ and Lundström K 1998b. Relationship between oxidation and conjugation metabolism of skatole in pig liver and concentrations of skatole in fat. *Journal of Animal Science* 76, 829–838.
- Babol J, Zamaratskaia G, Juneja RK and Lundström K 2004. The effect of age on distribution of skatole and indole levels in entire male pigs in four breeds: Yorkshire, Landrace, Hampshire and Duroc. *Meat Science* 67, 351–358.
- Barton-Gade PA 1987. Meat and fat quality in boars, castrates and gilts. *Livestock Production Science* 16, 187–196.
- Bee G 2001. Dietary conjugated linoleic acids affect tissue lipid composition but not de novo lipogenesis in finishing pigs. *Animal Research* 50, 383–399.
- Bee G, Gebert S and Messikommer R 2002. Effect of dietary energy supply and fat source on the fatty acid pattern of adipose and lean tissues and lipogenesis in the pig. *Journal of Animal Science* 80, 1564–1574.
- Bee G, Calderini M, Biolley C, Guex G, Herzog W and Lindemann MD 2007. Changes in the histochemical properties and meat quality traits of porcine muscles during the growing-finishing period as affected by feed restriction, slaughter age, or slaughter weight. *Journal of Animal Science* 85, 1030–1045.
- Bee G, Jacot S, Guex G and Biolley C 2008. Effects of two supplementation levels of linseed combined with CLA or tallow on meat quality traits and fatty acid profile of adipose and different muscle tissues in slaughter pigs. *Animal* 2, 800–811.
- Bergsma R, Knol E and Feitsma H 2007. Parameters of AI boars and predicted correlated responses of selection against boar taint. In 58th Annual Meeting of the European Association for Animal Production. Wageningen Academic Publishers, Dublin.
- Black JL, Giles LR, Wynn PC, Knowles AG, Kerr CA, Jones MR, Strom AD, Gallagher NL and Eamens GJ 2001. A review – factors limiting the performance of growing pigs in commercial environments. In *Manipulating pig production VIII* (ed. PD Cranwell), pp. 9–36. Australasian Pig Science Association, Werribee, Australia.
- Bonneau M 1982. Compounds responsible for boar taint, with special emphasis on androstenone: a review. *Livestock Production Science* 9, 687–705.
- Bonneau M 1998. Use of entire males for pig meat in the European Union. *Meat Science* 49(suppl. 1), S257–S272.
- Bonneau M, Denmat M, Vaudelet C, Veloso-Nunes JR, Mortensen AB and Mortensen HP 1992. Contribution of fat androstenone and skatole to boar taint I. Sensory attributes of fat and pork meat. *Livestock Production Science* 32, 63–80.
- Bonneau M, Kempster AJ, Claus R, Claudi-Magnussen C, Diestre A, Tornberg E, Walstra P, Chevillon P, Weiler U and Cook GL 2000. An international study on the importance of androstenone and skatole for boar taint: I. Presentation of the programme and measurement of boar taint compounds with different analytical procedures. *Meat Science* 54, 251–259.
- Campbell RG, Steele NC, Caperna TJ, McMurtry JP, Solomon MB and Mitchell AD 1989. Interrelationships between sex and exogenous growth hormone administration on performance, body composition and protein and fat accretion of growing pigs. *Journal of Animal Science* 67, 177–186.
- Chen G, Zamaratskaia G, Andersson HK and Lundström K 2007. Effects of raw potato starch and live weight on fat and plasma skatole, indole and androstenone levels measured by different methods in entire male pigs. *Food Chemistry* 101, 439–448.
- Claus R and Weiler U 1987. Umwelteinflüsse auf das geschlechtsspezifische Wachstumsvermögen. *Übersichten zur Tierernährung* 15, 301–316.
- Claus R, Weiler U and Herzog A 1994. Physiological aspects of androstenone and skatole formation in the boar: review with experimental data. *Meat Science* 38, 289–305.
- Claus R, Losel D, Lacorn M, Mentschel J and Schenkel H 2003. Effects of butyrate on apoptosis in the pig colon and its consequences for skatole formation and tissue accumulation. *Journal of Animal Science* 81, 239–248.
- Cronin GM, Dunshea FR, Butler KL, McCauley I, Barnett JL and Hemsworth PH 2003. The effects of immuno- and surgical-castration on the behaviour and consequently growth of group-housed, male finisher pigs. *Applied Animal Behaviour Science* 81, 111–126.
- D'Souza DN and Mullan BP 2002. The effect of genotype, sex and management strategy on the eating quality of pork. *Meat Science* 60, 95–101.
- Desmoulin B, Bonneau M, Frouin A and Bidard JB 1982. Consumer testing of pork and processed meat from boars: the influence of fat androstenone level. *Livestock Production Science* 9, 707–715.
- Diaz GJ and Squires EJ 2000. Metabolism of 3-methylindole by porcine liver microsomes: responsible cytochrome P450 enzymes. *Toxicological Sciences* 55, 284–292.
- Dunshea FR, King RH, Campbell RG, Sainz RD and Kim YS 1993. Interrelationships between sex and ractopamine on protein and lipid deposition in rapidly growing pigs. *Journal of Animal Science* 71, 2919–2930.
- Dunshea FR, Colantoni C, Howard K, McCauley I, Jackson P, Long KA, Lopaticki S, Nugent EA, Simons JA, Walker J and Hennessy DP 2001. Vaccination of boars with a GnRH vaccine (Improvac) eliminates boar taint and increases growth performance. *Journal of Animal Science* 79, 2524–2535.
- Fredriksen B, Lium BM, Marka CH, Heier BT, Dahl E, Choinski JU and Nafstad O 2006. Entire male pigs in a farrow-to-finish system. Effects on androstenone and skatole. *Livestock Science* 102, 146–154.
- Giersing M, Lundström K and Andersson A 2000. Social effects and boar taint: significance for production of slaughter boars (*Sus scrofa*). *Journal of Animal Science* 78, 296–305.
- Hansen-Moller J 1994. Rapid high-performance liquid chromatographic method for simultaneous determination of androstenone, skatole and indole in back fat from pigs. *Journal of Chromatography B. Biomedical Applications* 661, 219–230.
- Hansen LL, Larsen AE, Jensen BB, Hansen-Moller J and Barton-Gade PA 1994. Influence of stocking rate and faeces deposition in the pen at different temperatures on skatole concentration (boar taint) in subcutaneous fat. *Animal Production* 59, 99–110.
- Honikel KO 1998. Reference methods for the assessment of physical characteristics of meat. *Meat Science* 49, 447–457.
- Jensen MT, Cox RP and Jensen BB 1995. Microbial production of skatole in the hind gut of pigs given different diets and its relation to skatole deposition in backfat. *Animal Science* 61, 293–304.
- Lin ZH, Lou YP and Squires EJ 2006. Functional polymorphism in porcine CYP2E1 gene: Its association with skatole levels. *Journal of Steroid Biochemistry and Molecular Biology* 99, 231–237.
- Lösel D 2006. Versuche zur Verbesserung der sensorischen Fleischqualität beim Schwein durch nutritive Hemmung der Skatolbildung. Universität Hohenheim.
- Lösel D and Claus R 2005. Dose-dependent effects of resistant potato starch in the diet on intestinal skatole formation and adipose tissue accumulation in the pig. *Journal of Veterinary Medicine A. Physiology, Pathology, Clinical Medicine* 52, 209–212.
- Malmgren L 1993. Induced testicular alterations in prepubertal and mature boars. Thesis. Department of Obstetrics and Gynaecology, Swedish University of Agricultural Sciences, Uppsala.
- Martin LJM, Dumon HJW, Lecanu G and Champ MMJ 2000. Potato and high-amylose maize starches are not equivalent producers of butyrate for the colonic mucosa. *British Journal of Nutrition* 84, 689–696.
- Mentschel J and Claus R 2003. Increased butyrate formation in the pig colon by feeding raw potato starch leads to a reduction of colonocyte apoptosis and a shift to the stem cell compartment. *Metabolism* 52, 1400–1405.
- Morano I, Gerstner J, Ruegg JC, Ganten U, Ganten D and Vosberg HP 1990. Regulation of myosin heavy chain expression in the hearts of hypertensive rats by testosterone. *Circulation Research* 66, 1585–1590.
- Pauly C, O'Doherty JV and Spring P 2007. Performance and carcass quality of entire male pigs fattened on a commercial farm in Switzerland. In 58th Annual Meeting of the European Association for Animal Production, Dublin.
- Prunier A, Bonneau M, von Borell EH, Cinotti S, Gunn M, Fredriksen B, Giersing M, Morton DB, Tuytens FAM and Velarde A 2006. A review of the welfare consequences of surgical castration in piglets and the evaluation of non-surgical methods. *Animal Welfare* 15, 277–289.
- Rydhmer L, Zamaratskaia G, Andersson HK, Algers B, Guillemet R and Lundström K 2006. Aggressive and sexual behaviour of growing and finishing pigs reared in groups, without castration. *Acta Agriculturae Scandinavica Section A. Animal Science* 56, 109–119.
- Sather AS, Jeremiah LE and Squires EJ 1999. The effects of castration on live performance, carcass yield and meat quality of male pigs fed wheat or corn-based diets. *Journal of Muscle Food* 10, 245–259.

- Scheeder MRL, Bossi H and Wenk C 1999. Kritische Betrachtungen zur Fettzahl-Bestimmung. *Agrarforschung* 6, 1–8.
- Schnegg HU, Stoll P and Jost M 1985. Häufigkeit von Ebergeruch bei der Schlachtung von Jungebern unterschiedlicher Herkunft. In *Interner Versuchsbericht der FAG*.
- Schoonmaker JP, Loerch SC, Fluharty FL, Turner TB, Moeller SJ, Rossi JE, Dayton WR, Hathaway MR and Wulf DM 2002. Effect of an accelerated finishing program on performance, carcass characteristics, and circulating insulin-like growth factor I concentration of early-weaned bulls and steers. *Journal of Animal Science* 80, 900–910.
- Stoll P 1982. Voraussetzungen für die Mast von Jungebern: Aspekte der Fleischbeschaffenheit und der Geruchsabweichungen. In *Fleischproduktion mit unkastrierten männlichen Schweinen*, Eidgenössische Forschungsanstalt für viehwirtschaftliche Produktion, Posieux.
- Suster D, Leury BJ, Kerton DJ, Borg MR, Butler KL and Dunshea FR 2006. Longitudinal DXA measurements demonstrate lifetime differences in lean and fat tissue deposition between boars and barrows under individual and group-penned systems. *Australian Journal of Agricultural Research* 57, 1009–1015.
- Turkstra JA, Zeng XY, van Diepen JTM, Jongbloed AW, Oonk HB, van de Wiel DFM and Meloen RH 2002. Performance of male pigs immunized against GnRH is related to the time of onset of biological response. *Journal of Animal Science* 80, 2953–2959.
- Walstra P, Claudi-Magnussen C, Chevillon P, von Seth G, Diestre A, Matthews KR, Homer DB and Bonneau M 1999. An international study on the importance of androstenone and skatole for boar taint: levels of androstenone and skatole by country and season. *Livestock Production Science* 62, 15–28.
- Whittemore CT, Tullis JB and Emmans GC 1988. Protein growth in pigs. *Animal Production* 46, 437–445.
- Xue J, Dial GD and Pettigrew JE 1997. Performance, carcass, and meat quality advantages of boars over barrows: a literature review. *Swine Health and Production* 5, 21–28.
- Zamaratskaia G, Babol J, Andersson H and Lundström K 2004. Plasma skatole and androstenone levels in entire male pigs and relationship between boar taint compounds, sex steroids and thyroxine at various ages. *Livestock Production Science* 87, 91–98.
- Zamaratskaia G, Babol J, Andersson HK, Andersson K and Lundström K 2005a. Effect of live weight and dietary supplement of raw potato starch on the levels of skatole, androstenone, testosterone and oestrone sulphate in entire male pigs. *Livestock Production Science* 93, 235–243.
- Zamaratskaia G, Rydhmer L, Chen G, Madej A, Andersson HK and Lundström K 2005b. Boar taint is related to endocrine and anatomical changes at puberty but not to aggressive behaviour in entire male pigs. *Reproduction in Domestic Animals* 40, 500–506.
- Zeng XY, Turkstra JA, Jongbloed AW, van Diepen JTM, Meloen RH, Oonk HB, Guo DZ and van de Wiel DFM 2002. Performance and hormone levels of immunocastrated, surgically castrated and intact male pigs fed ad libitum high- and low-energy diets. *Livestock Production Science* 77, 1–11.