

Heritability and phenotypic correlations of protein efficiency in a Swiss Large White pig population

C. Kasper¹, I. Ruiz-Ascacibar¹, P. Stoll¹ and G. Bee¹

¹Agroscope, La Tioleyre 4, 1725 Posieux, Switzerland

Introduction Pig production contributes importantly to environmental pollution through the emission of nitrogen waste. Many countries cannot meet the plant-protein demand of livestock with current feed cultivation practices, and an important part of high-quality protein feed, most importantly soybean meal, is imported from South America. The high global demand for protein results in an increased competition of agricultural land for human food and animal feed and ultimately to the large-scale loss of ecological habitats. It is therefore desirable to improve protein efficiency, i.e. the proportion of dietary protein that is fixed in the body, of pigs, an important livestock species, through selective breeding. However, little is known about the inheritance pattern and genetic architecture of protein efficiency as well as the genes underlying protein efficiency and their functions, which is essential for assessing the potential to breed pigs with higher protein efficiency.

Materials and Methods We used data of 294 offspring of 17 sires and 56 dams from previous experiments (Ruiz-Ascacibar et al., 2017). Nitrogen (~protein) efficiency of the empty body (NE_{effEB}) and of the carcass ($NE_{effCarc}$) were the proportion of fixed protein in the entire body and the carcass, respectively, from the total dietary protein intake. We estimated the genetic (h^2) and environmental variance components (CE^2) of NE_{effEB} and $NE_{effCarc}$ using a mixed-effect Gaussian animal model (Henderson, 1984) following a Bayesian framework (Hadfield, 2010). We assessed phenotypic correlations of NE_{effEB} with phosphorus efficiency, water, raw ash and fat content of the empty body of a subset (N=73) for which this information was available. The correlation of NE_{effEB} with the number of days slaughtering was delayed because a pig did not reach the target body weight was calculated for the entire dataset.

Results The heritability of N efficiency of the empty body was higher than the one of the carcass ($h^2_{NE_{effEB}} = 0.32 [0.15, 0.59]$ and $h^2_{NE_{effCarc}} = 0.16 [0.08, 0.41]$) (Fig. 1). The common environment also contributed to the phenotypic variation in NE_{effEB} ($CE^2 = 0.13 [0.08, 0.27]$) and $NE_{effCarc}$ ($CE^2 = 0.16 [0.09, 0.25]$). We found a low negative phenotypic correlation of NE_{effEB} and the number of days an animal was delayed in reaching target weight (Table 1). NE_{effEB} correlated moderately positively with phosphorus efficiency. Body composition traits (water/crude ash/crude fat content) were not correlated with NE_{effEB} .




Table 1 Correlations of NE_{effEB} with traits of potential economic and ecological impact as well as with body composition.

trait	type	coefficient \pm 95% CI
days delayed	Spearman	$\rho = -0.16 [-0.26, -0.07]$
P efficiency	Pearson	$r = 0.65 [0.48, 0.77]$
water content	Pearson	$r = -0.07 [-0.32, 0.18]$
crude ash content	Pearson	$r = -0.16 [-0.39, 0.09]$
crude fat content	Pearson	$r = -0.14 [-0.38, 0.11]$

Figure 1 Heritability (h^2 , blue), common environment effect (CE^2 , green) and residual variance (r^2 , red) of protein efficiency of the whole body (A) and protein efficiency of the carcass (B). Posterior distributions of the respective variance components (upper part), points representing single estimates are shown together with a box plot (with median, interquartile range and 5th to 95th percentile range).

Conclusions Our preliminary results indicate a potential for selective breeding towards increased protein efficiency but more research is needed to improve estimates of genetic parameters. Information on the association of genetic loci (SNPs) and protein efficiency will help understand the genes and their functions that underlie protein efficiency. We found no major trade-offs with other economically important traits. Breeding for increased protein efficiency could lead to a slower growth of the animals, but the decrease is expected to be small and might be offset by lower costs of protein-reduced feed. However, potential trade-offs with fertility, aggression and meat quality should be addressed by future studies. A reduction of proteins in pig feed can potentially facilitate both conventional and organic pig production in Switzerland.

References

- Hadfield, J.D., 2010. MCMC Methods for Multi-Response Generalized Linear Mixed Models: The **MCMCglmm** R Package. *J. Stat. Softw.* 33. <https://doi.org/10.18637/jss.v033.i02>
- Henderson C.R., 1984. Applications of Linear Models in Animal Breeding. University of Guelph, Guelph, CA, 462 pages
- Ruiz-Ascacibar, I., Stoll, P., Kreuzer, M., Boillat, V., Spring, P., Bee, G., 2017. Impact of amino acid and CP restriction from 20 to 140 kg BW on performance and dynamics in empty body protein and lipid deposition of entire male, castrated and female pigs. *animal* 11, 394–404. <https://doi.org/10.1017/S1751731116001634>