



Contents lists available at SciOpen

Food Science and Human Wellness

journal homepage: <https://www.sciopen.com/journal/2097-0765>

Tailored Co-Extruded Cereals for Seniors: Fabrication, Palatability, and *in vitro* Digestibility

Leehen Mashiah ^a, Omer Medini ^b, Andrea Araiza Calahorra ^c, Josefine Skaret ^d, Carmit Shani Levi ^a,
Lotti Egger ^c, Reto Portmann ^c, Paula Varela ^d, Anwasha Sarkar ^e, Uri Lesmes ^{a,*}

^a Laboratory of Chemistry of Foods and Bioactives, Department of Biotechnology and Food Engineering, Technion – Israel Institute of Technology, Haifa, Israel

^b The International Beer Breweries LTD, Kibbutz, Givat Haim, M.P. Hefer, 3893000, Israel

^c Agroscope, Schwarzenburgstr, 161, 3003 Bern, Switzerland

^d Nofima AS. P.O. Box 210, 1431 Ås, Norway

^e Food Colloids and Bioprocessing Group, School of Food Science and Nutrition, University of Leeds, Leeds, LS2 9JT, UK

ABSTRACT: The rise in life expectancies and gap between life span and health span necessitate innovative approaches, interventions, and food solutions to secure healthy aging. This study, conducted within the framework of the EAT4AGE project of the Joint Programming Initiative 'A Healthy Diet for a Healthy Life', focuses on the rational design of functional foods for older adults. A co-extruded cereal prototype, fortified with two bioactive moieties, Maca root powder and Olive leaf extract, aims to address nutritional gaps identified in older adults, with a particular emphasis on high-quality and highly digestible proteins. Analytical determinations reveal the cereals have rich macronutrient profile exceeding 12% (w/w) protein, 20% (w/w) fat, and low sugar (< 5% w/w), surpassing commercially available products with texture analyses supporting improved hardness, reduced oral friction and oral comfort. An untrained consumer panel (n=21, Age 73 ± 5) confirmed high palatability in various metrics and overall acceptability that were also affirmed through a trained sensory panel. Lastly, the product digestion was explored through an age-tailored *in vitro* digestion model which consistently demonstrated high protein digestibility, surpassing 80%, across all product formulations. Further, calculation of the *in vitro* digestible indispensable amino acid score of the product affirms its high nutritional quality. Thus, this study underscores the potential of designing palatable foods that could help promote a balanced and sustainable diet towards healthy aging.

Keywords: Healthy aging, functional foods, co-extrusion, palatability, oral comfort, plant-based cereals, *In vitro* protein digestibility, DIAAS, proteomic analysis, breakfast cereals

1. Introduction

The National Institutes of Health (NIH), U.S identified a pronounced rise in human lifespan that is yet to be matched by a concomitant rise in human health span [1], [2]. This is accompanied by a rise in efforts to exhaust the potential of plant-based foodstuffs towards healthier and more sustainable diets [3], [4]. Thus, development of tailored nutritional solutions can significantly affect the independence, quality of life and

*Corresponding author
lesmesu@bfe.technion.ac.il

Received 5 February 2024
Received in revised form 26 March 2024
Accepted 26 August 2024

healthy aging of older adults by addressing many of the adversities of aging [5], [6]. Such solutions need to address relevant age-related physiological decline, e.g. changes in salivation, reduced mandibular forces, problems swallowing, slower gastric emptying, decreased excretion of digestive fluids and altered microbiota along with decreased appetite and early satiety [5], [7], [8], [9], [10]. In addition, design of foods for the older adults' mandates tackling age-related mental decline and changes in eating patterns, wherein potential sensory deterioration may underlie malnourishment and insufficient protein intake and damper health [5], [11], [12], [13].

Today, there is a growing consumer and commercial interest in food and dietary solutions that can effectively meet the unique preferences, acceptance and needs of seniors [5], [12]. This led to an ERA-Net ERA-HDHL call for the "Development of targeted nutrition for prevention of undernutrition for older adults" (PREVNUT). This challenge was met by several consortia, one of which was the EAT4AGE consortium that targeted the development of palatable nutrient-dense functional foods. Within this project, a literature survey pin-pointed actual nutritional gaps in older adults, with high quality protein, dietary fibres and specific micronutrients, like calcium and iron set as viable targets [11], [14], [15], [16], [17]. In fact, protein was found to be of great nutritional importance, as a robust dietary interventional tool to affect sarcopenia of aging, healthy muscle ageing and even help regulate appetite [17], [18], [19], [20]. To this end, senior consumption of dietary proteins has been suggested to be increased to 1.0-1.2 gr/kgBW/day (compared to 0.8 gr/kgBW/day recommendation for healthy adults) with indications that animal proteins, e.g. dairy proteins, exhibiting improved digestibility and nutritional value in older adults compared to plant-based protein sources, e.g. soy [16], [17]. However, animal proteins are not devoid of limitations and plant-based proteins prove to be advantageous in various aspects, notably as part of recommended dietary choices leaning towards healthier and eco-friendlier planetary diets [3], [4], [21], [22].

In addition, this work sought to offer seniors with value-added products by incorporating functional food bioactives [23], [24], [25]. Out of the various possibilities, this work identified and focused on Macamides found in the Maca root (Maca) and Oleuropein in Olive Leaf Extract (OLE) [26], [27], [28]. Maca was selected for its various indications of beneficial antioxidant, hormone-balancing and therapeutic properties that are relevant to aged individuals, these include enhanced fertility [29] and alleviation of symptoms associated with menopause [30]. Supplementation with OLE was selected due to demonstrated safety [31] and indications that its supplementation reduces mitochondrial superoxide production and increases mitochondrial biogenesis[32], contributing to an overall positive impact on cellular health.

Applying consumer-oriented considerations for the development of functional foods for seniors gave rise to a pragmatic solution in the form of a shelf-stable extruded cereal product. This relied on indications that consumption of extruded cereal products and plant-based products has increased over the years due to their considerable shelf life, affordability, sustainability, and ease of use as ready-to-eat foods [33], [34]. Moreover, manufacturing of such plant-based products, produced via extrusion, may help tackle challenges of allergens, antinutritional factors (ANF) and improve their overall limited digestibility [35], [36], [37]. For instance,

extrusion process has been shown to promote denaturation of proteins and inactivation of thermo-labile ANF moieties, thereby enhancing *in vitro* protein digestibility and positively influencing protein quality [36], [37].

This rational food engineering approach sought to open opportunities for tailoring food compositions to tackle the challenges of maximizing palatability, oral comfort, lubricity while optimizing nutrient bioaccessibility and bioavailability to seniors [38], [39], [40], [41]. In fact, a growing number of recent studies demonstrate that the altered digestive functions of seniors affect their ability to render nutrients bioaccessible [5], [42], [43], [44], [45]. Furthermore, interactions between food components may interfere with nutrient digestibility and absorption [46], [47], [48]. Consequently, research into the digestive fate of tailored food products is an essential milestone towards their bio-efficacious development [5], [40], [44].

Thus, this research sought to design and study the potential digestive fate of a co-extruded cereal product in seniors (age > 65). Moreover, the product was formulated to be a functional food with two added bioactive moieties: Maca and OLE from the olive oil industry. This study focused on the palatability of the functional product, its oral processing and the potential digestibility of its proteins in healthy seniors. The underlying hypothesis was that seniors have altered food preferences and digestive capabilities which affect their perception and breakdown of the tailored cereal product. For this purpose, we adapted a two-armed study combining *in vitro* and *in vivo* measures.

2. Materials and Methods

To investigate the potential digestive fate of the functional cereals in healthy older adults, we employed a battery of methods delving into the oral processing of the product and its gastro-intestinal digestibility using and *in vitro* setup, guided by a recently published consensus protocol [9].

2.1. Materials

Chickpea, rice and teff flours, coconut oil and tapioca starch were acquired in a local supermarket (Shufersal Ltd. Israel). Olive Leaf Extract (OLE) with 20% oleuropein (Doctor Phyto, Israel) was acquired from Bio-Gaya (Bio-Gaya, Israel). Maca powder (Beit Shaked Kfar, Israel) was acquired in a local supermarket. Roasted Sesame bran and sesame paste (tahini) were generously donated by Al-Arz (Al Arz, Israel). Magnesium carbonate, corn grits, cocoa, saccharin, vanillin, NUTRIOSE and erythritol were kindly provided by the International Beer Breweries LTD pilot laboratory. Fruitlift™ is an all-natural sweetening solution containing 90% fruit components, fully compatible with extrusion processing (ensuring no “caking” and no stickiness). All the above ingredients were food grade, and used as received without any purification.

Standard 1000 ppm Single Element Standards by Agilent technologies mineral solutions for mineral composition analysis was used. Pepsin from porcine gastric mucosa (P7000); Trypsin from porcine pancreas (T0303); α -Chymotrypsin from bovine pancreas (C4129); Pancreatin from porcine pancreas (P7545); All enzymes batches were studied for their specific activity (including periodical tests to monitor changes during shelf life) and used according to the INFOGEST protocol and its overall rationale [49]. Porcine bile extract (SC-214601) was purchased from Santa Cruz Biotechnology. Sodium glycodeoxycholate; taurocholic acid sodium salt hydrate; phenylmethylsulfonyl fluoride (PMSF); pepstatin-A (CAS# 26305-03-3, microbial,

≥90% (HPLC)) were purchased from Sigma Aldrich (Rehovot, Israel). Calcium chloride dehydrate was obtained from Spectrum Chemical Manufacturing Corporation (New Brunswick, NJ, USA). All chemicals and enzymes used in the present study for determination of total amino acid were purchased from MERCK (Zug, Switzerland). Gels were stained by Instant blue stain (Bio Consult, Israel). All other chemicals and reagents used in the study were of analytical grade.

2.2. Formulation and co-extrusion of cereal products

Acknowledging "anabolic resistance" in older adults [5], the product was designed to offer easily digestible, high-quality proteins in a shelf-stable and accessible food product. This relied on studies showing protein bioaccessibility and digestibility of plant proteins are improved by their controlled processing [36], [50], [37], [38]. Thus, a calorie-dense product was crafted to comprise a shell made from five gluten-free flours that also offer a well-balanced AA profile along with sesame peels, a by-product of tahini production and a rich source of dietary fibers[51]. To enhance functionality, the product was also supplemented with two key bioactive moieties whose bioactivity is supported by clinical studies, Maca was added to the shell and OLE added to the oily filling [27], [30], [32], [52], [53]. The first additive, Maca powder, was selected for its various beneficial indications relevant to hormonal regulation and immunomodulation in seniors [54], [55]. Similarly, OLE was selected due to indications that its supplementation has antidiabetic and antioxidant effects as well as beneficial effects on mitochondrial function [32], [56].

The complex hard shell comprised teff, chickpea, maize, tapioca, and rice with Fruitlift™ for sweetening and Maca root powder as functional ingredient. The filling was a mixture of sesame paste, coconut oil, flavorings (vanillin, cocoa powder, saccharin, erythritol) and OLE as a functional supplement. The production of three test products was pursued through co-extrusion at International Beer Breweries Ltd. that enabled the production and packaging of the food-grade ready-to-eat products (see formulae in **Table 1**). Co-extrusion of the plant-based cereal product was optimized for production and three formulae were produced: a control or reference version without any functional ingredients in the shell or the filling (Ref+ref), a product with 0.03% w/w Maca-added to the shell (Maca+ref) and a product with 0.03% w/w Maca and 0.015% w/w OLE in the shell and filling (Maca+OLE), respectively.

The products were produced via co-extrusion using a 36 mm twin-screw extruder (L/D=28, main die 6 mm diameter and core filling nozzle diameter 4 mm) in a pilot plant complying with food-grade national health and safety regulations. The screw elements, including kneading blocks and reverse screw elements were selected to offer high shear at 220 r/min. Feed was conveyed into the extruder with a twin-screw weight feeder (Shandong Light M&E Co) and adjusted to a feed rate of 10 kg/hr. The die temperature was 105°C and samples were collected after 3 min of processing. The samples were air dried on trays at room temperature, then approximately 15gr of product were packed in aluminium bags sealed with heat welding of the polyethylene component. All samples were stored at room temperature until further analysis.

Table 1. Composition of co-extruded plant-based cereal products

Ingredients (amount in 1 kg product)		Ref+ref	Maca+ref	Maca+oleu
Shell	Corn grits (gr)	360	340	340
	Chickpea (gr)	200	200	200
	Rice (gr)	110	100	100
	Tapioca (gr)	100	100	100
	FruitLift solution (gr)	100	100	100
	Teff (gr)	80	80	80
	Maca root powder (gr)	-	50	50
	Sesame peels	20	20	20
	Cocoa (gr)	10	10	10
	Salt (gr)	3	3	3
Filling	Magnesium carbonate (gr)	3	3	3
	Tahini (gr)	600	600	600
	Erythritol (gr)	150	150	150
	NUTRIOSE (gr)	150	150	150
	Coconut oil (gr)	80	80	80
	Cacao (gr)	20	20	20
	Vanillin (gr)	0.3	0.3	0.3
	Saccharin (ppm)	40	40	40
	Olive leaf extract (ml)	-	-	15

*The pivotal process parameters for co-extrusion encompassed inlet and outlet temperatures of 13°C and 100/105°C, respectively, coupled with a pressure of 553 mPa, a die diameter of 6 mm (L/D=28) in a 36 mm twin-screw extruder. Initial moisture content set at 13%-15% (w/w) and a final target of 7.2% (w/w).

2.3. In-vitro characterization of the plant-based cereal prototypes

2.3.1. Proximate analyses

Protein content was determined by the Kjeldahl method and by automated CHNS analysis system enabling DUMAS protein quantification. CHNS Elemental analysis was performed using Flash 2000 Organic elemental analyzer (Thermo Scientific, USA). Samples of 2-4 mg were weighed with 8-10 mg of vanadium in a tin crucible. Combustion Temp. was 950°C, carrier gas was He (99.999%); 140 ml/min. Adding O₂ at 250 ml/min for 5 seconds. Standards: Cystine, BBOT, Sulphanilamide, Methionine, Nicotinamide. A conversion factor of 5.7 was used for the cereal products [57]. The fat content was determined according to AOAC method 996.01[58]. All analyses were performed in triplicate. Regulatory labeling values were calculated based on individual ingredient composition and customized software analysis pack (Tzameret software, Department of Nutrition, Israel Ministry of Health).

2.3.2. Instrumental single compression test

Instrumental single compression tests were carried out using a TA1 texture profile analyzer (Lloyd Instruments Ltd, AMETEK, Israel) equipped with a cylindrical plate (115 mm diameter, working force 50 N, pre-speed and test speed set to 3 mm/s and 1 mm/s, respectively)[59], [60], [61]. A single sample of each of the three products was placed on the plate and its deformation was observed for a compression up to 90% of the original product thickness[59]. The parameters hardness (N) and Young's modulus were obtained from the instrumental single compression tests[62].

2.3.3. Tribological measurements

The evaluation of the lubrication properties of the simulated bolus samples was conducted at a controlled temperature of 37°C using a Mini-Traction Machine (MTM2) provided by PCS Instruments, UK. This equipment offers the advantage of measuring the friction coefficient across a broad speed range, enabling a comprehensive mapping of the Stribeck curve. The tribopair configuration involved a ball (19.0 mm diameter) on disc contact, where both surfaces were composed of polydimethylsiloxane (PDMS) with a Young's modulus of 2.4 MPa and an average surface roughness (Ra) of approximately 50 nm [63]. Prior to the experiments, the PDMS surfaces were cleaned in an ultrasonic bath with a solution of 3% surface-active cleaning agent (Decon 90®, East Sussex, UK), a solution of 10% isopropanol, followed by rinsing with isopropanol. After such treatment, the surface of the PDMS retained its natural hydrophobic characteristic.

For measurement, pure sliding conditions of 50% slide-roll ratio (SRR) was used and three tests of ascending sliding speed (0.1–1,000 mm s⁻¹) were completed, and the average reported. A volume-reducing insert was used allowing for a sample size of 15 mL. Experiments were performed at 37 °C. A normal force of 2 N was used as in-mouth friction have been reported to be between 0.1 N and 10 N [64]. Hence, 2 N would allow low contact pressure, which is of relevance in oral processing applications.

2.3.4. Microstructure characterisation by Confocal scanning laser microscopy (CLSM)

The microstructures of the simulated sample boli were characterized before being subjected to tribological stress using a Zeiss LSM 880 inverted confocal microscope (Carl Zeiss MicroImaging GmbH, Jena, Germany). Approximately 5 µL of Nile Red solution (0.1% v/w in dimethyl sulfoxide) and Fast Green solution (1.0% v/w in MilliQ water) were used to fluorescently stain the lipid (excited at 514 nm) and protein phases (excited at 633 nm), respectively. The fluorescently labeled sample boli were placed on a concave confocal microscope slide, fixed with a glass coverslip, and imaged using an oil-immersion 63 × lens and pinhole diameter of 1 Airy Unit to filter out most of the light scattering.

2.3.5. Simulated oral processing

2.3.5.1. Preparation of model saliva

The model saliva was prepared following the composition previously described by [65]. Briefly, to prepare 1 L of model saliva, 1.59 g L⁻¹ NaCl, 0.328 g L⁻¹ NH₄NO₃, 0.64 g L⁻¹ KH₂PO₄, 0.20 g L⁻¹ KCl, 0.31 g L⁻¹ K₃C₆H₅O₇·H₂O, 0.02 g L⁻¹ C₅H₃N₄O₃Na, 0.20 g L⁻¹ H₂NCONH₂, 0.15 g L⁻¹ C₃H₅O₃Na and 3.00 g L⁻¹ porcine gastric mucin type II were dissolved in distilled water. After adjusting the pH to 7.0 using 1 M NaOH, the volume was made up to 1 L using a volumetric flask. Porcine gastric mucin was used due to its ability to simulate the rheological properties of human saliva. It is noteworthy, however, that bovine submaxillary mucin is the optimal source of commercially available mucin for lubricating properties[66], and therefore this is a limitation of the current study.

2.3.5.2. Simulated bolus formation

To simulate oral processing, food boli were prepared as described in Menard et al., 2023[9]. Briefly, samples were mixed with model saliva at a final insalivation ratio, food: saliva ratio of 1:1, which was

determined from spit boli obtained from INRAE. Subsequently, the samples underwent mincing using a Kitchen Craft No. 5 meat mincer equipped with a 5 cm mincing disc and a 0.5 cm mesh size for one pass. After recovering the bolus, additional model saliva was introduced to achieve a final insalivation ratio of 1:4. The mixture was agitated using a magnetic stirrer at 100 r/min for 10 minutes to ensure particle dispersion while preserving the bolus structure, thereby achieving a consistency suitable for subsequent characterization.

2.4. Comprehensive in vivo sensory evaluation

2.4.1. Food comfortability

Food comfortability tests were evaluated by 21 untrained volunteers (mean age: 73 ± 5 , 13 females, 8 males), with specific criteria established to ensure a representative sample of older adults. This study cohort was recruited to meet levels for laboratory acceptability testing [67] and using a previously developed questionnaire [68]. Participants were recruited from the community and each participant gave written consent to participate in the experiments as well as a declaration that he/she complied with the study's inclusion criteria (over 65, overall independent community-dwelling, healthy and in good present state, allergy-free, and not suffering from dry mouth or other oral or gastrointestinal problems). The subjects who met all the requirements were invited to an information session, where they received an explanation of the study setup and signed the informed consent form. During the sessions, the volunteers were invited to eat a portion of the three formulae cereal products prior to each of the 5 parts of the questionnaire (water was available ad libitum). The evaluations were performed under normal light at room temperature (approximately 25 °C).

2.4.2. Sensory evaluation of products

Ten experienced assessors, highly trained in the use of quantitative descriptive analysis (QDA) and Temporal Dominance of Sensations (TDS) participated in this part of the research. The formal assessments were conducted in individual booths under white light with proper ventilation, while general discussions among panelists were conducted in a well-ventilated discussion room. The sensory laboratory was designed according to guidelines in ISO 8589:2007(E) featuring separate booths and electronic registration of data (Eye Question, v. 3.8.6, Logic 8, The Netherlands), standardized lighting and a separate ventilation system.

2.4.2.1. Quantitative Descriptive Analysis (QDA)

Nofima's trained panel conducted the sensory profiling on the cereal samples using QDA as described previously [69]. The vocabulary for the products was established during a preliminary session involving two of the samples. This pre-trial session, lasting one hour, involved the assessors in agreeing upon descriptors, their definitions, and reference samples. By the end of the pre-trial, all assessors demonstrated the ability to distinguish among samples, showed consistency in their evaluations during subsequent trials, and reached a consensus within the group. The final list of attributes included odor, flavor, texture and mouthfeel attributes: Sweetly odor, Chocolate odor, Nutty odor, Roasted odor, Grain odor, Drawer odor, Rancid odor, Sweet taste, Salty taste, Bitter taste, Chocolate flavor, Nutty flavor, Roasted flavor, Grain flavor, Drawer flavor, Cloying flavor, Rancid flavor, Crispness, Juiciness, Fatness, Stickiness and Astringency.

Each panelist received each cereal sample in plastic cups labeled with randomly assigned 3-digit codes. The samples were presented at room temperature in a sequential monadic fashion following a balanced presentation order. The evaluations were conducted in two replicates and lasted approximately 1.5 hours.

2.4.2.2. Temporal dominance of sensations (TDS)

The assessment process followed the TDS methodology outlined previously [70]. TDS allows for the dynamic sensory description of food products, during the whole consumption period. Initially, assessors were reminded of the concept of dominant sensations during food consumption. Subsequently, they tasted the samples and noted the attributes they perceived to differentiate the samples from a temporal perspective (during the whole consumption period). The most relevant attributes for the dynamic description of the products were selected through discussion and consensus among the panelists.

The sensory lexicon created for the cereal samples included eight flavor and texture attributes with definitions sourced from ISO 5492:2008 (bitter, chocolate, roasted, grain, drawer, rancid, crispy, sticky). In the formal assessment, assessors evaluated the samples one by one, served in small plastic cups labeled with unique 3-digit codes. Water rinses were required between samples.

Assessors were instructed to place the sample in their mouth, press the "START" button, and choose the dominant sensations they experienced while consuming the sample by clicking on one of the eight attributes displayed on screen. When they were ready to swallow, they pressed "STOP" and spat out the sample. Assessors could choose from among the attributes as many times as necessary during the consumption of the samples, even re-selecting an attribute more than once during the test; only one attribute at a time could be selected as the dominant sensation.

2.5. *In vitro* semi-dynamic digestion adjusted to older adults

Since data on oral processing in older adults is scant [9], this part of the work initiated with estimating the salivary mineral composition of seniors ($n=10$, Age= 75 ± 5) by Inductively Coupled Plasma Optical Emission Spectroscopy (ICP-OES) with outcomes given in the supplementary materials (see **Table 1** in the supplementary materials). This affirmed that the calcium composition of seniors doesn't differ from that of young adults [49], validating that defined recently for *in vitro* digestion modeling [9]. The bolus features, however, can be impacted by other aging-related changes in oral processing [5], [71]. For this reason, oral bolus from senior volunteers ($n=4$) was selected to be used in the semi-dynamic *in vitro* digestion (IVD) model.

Samples were orally processed *in vivo* (four healthy volunteers, mean age= 75 ± 2) right before being subjected to semi-dynamic IVD. Each volunteer chewed a sample of 15 gr until a point where they felt ready to swallow but instead the oral bolus was spit it into a sterile cup. Subsequently, a gargle with 15 mL of water was performed to dislodge any remaining residue adhering to the teeth. As controls, the pre-extruded product blend (Mix, Mix+maca) or a protein-free cookie (comprised solely of fat and carbohydrates) were mixed in a 1:1 ratio with SSF solution and 75 μ L CaCl₂ (0.3 M) instead of being chewed by the volunteers.

Semi-dynamic *in vitro* digestion. The simulation of human gastrointestinal digestion was carried out following an adjusted form that combines two standardized methods for simulated IVD [49], [72], similar to that depicted in past studies [73], [74].

Simulated adult digestion. Measured oral bolus aliquots (30 mL) were placed in a stirred double jacketed vessel (6.1418.250, Metrohm, Switzerland) ($37 \pm 1^\circ\text{C}$ and ~ 250 r/min) which was controlled by a dual auto titration unit (Titrando 902, Metrohm, Switzerland). Then, 28 mL preheated simulated gastric fluid (SGF) (pH 3, 37°C) were added, followed by the addition of 2 mL freshly prepared pepsin solution (final concentration of 2000 U/mL) and 15 μL CaCl_2 (0.3 M). Once mixed in the vessel, the TIAMO control software was activated to follow a semi-dynamic digestion protocol and generate a 2 h gastric pH gradient (using 0.3 M HCl), as previously reported [75]. At the end of this gastric phase, half of the gastric digesta was collected and mixed with simulated intestinal fluid (SIF) (1:1 vol ratio). Trypsin (100 U/mL), alpha-chymotrypsin (25 U/mL), pancreatic alpha-amylase (200 U/mL), sodium glycodeoxycholate and taurocholic acid sodium salt hydrate (final concentration of 10 mmol/L in total) and 60 μL CaCl_2 (0.3 M) were added, and the intestinal phase protocol was initiated. This phase maintained at pH = 6.25 (using 0.3 M NaOH) for 2 h at (37 ± 1) $^\circ\text{C}$. A total of ten aliquots of 1 mL were aspirated during the digestion experiments: six from the gastric phase at time lapses of 0, 15, 30, 60 and 120 min (abbreviated as G0, G15, G30, G60 and G120); four samples from the intestinal phase at time lapses of 15, 30, 60 and 120 min (abbreviated as D15, D30, D60 and D120). Upon collection, each aspirate was inactivated: gastric effluents by pepstatin-A (final concentration of 7 $\mu\text{mol/L}$) and intestinal effluents using the irreversible serine-protease inhibitor PMSF (final concentration of 0.5 mmol/L). All inactivated gastric and intestinal effluents were stored at -20°C until further analyses. The pre-extruded product (Mix, Mix+maca) and a protein-free cookie (with only fat and carbohydrates) were digested in triplicates, as controls.

Simulated older adult digestion. Semi-dynamic IVD was performed similarly to adult digestion with modifications to mirror age-related physiological changes in gut functions, as detailed previously [5], [9], [42], [76]. Each product sample was subjected to two different sets of experiments: one set included three replicate experiments done using oral contributions from a single individual, and another set of digestion experiments with three contributions collected from three different individuals. All oral boli were subjected to a semi-dynamic gastro-intestinal protocol concurring with a recent consensus protocol devised for older adults [9]. Briefly, this model included changes in the pH and duration of the gastric phase, as well as the activity of the digestive enzymes in the stomach (pepsin with final concentration of 1200 U/mL) and small intestine (20% decrease for all enzymes) and the concentration of bile salts (6.7 mmol/L in total) and CaCl_2 in the intestinal (final concentration of 1 mmol/L). The composition of SSF, SGF, SIF concentrations were based on [42], [76] and detailed in **Table 2**. In the gastric phase, the pH gradient was adjusted to slightly different profile to stimulate the elevated gastric pH (major steps: pH=5 for 23 min, pH=4 for 52 min, pH=3 for 89 min, pH=2 for 16 min)[9]. The gastric phase was carried out for 3 h and the intestinal phase was kept at constant pH = 6.25 for

2 h. The schematic depiction (**Figure 1**) elucidates and defines the specific digestive conditions for seniors, offering a valuable framework for future research endeavours in this field.




Figure 1. Schematic illustration of simulated senior digestive tract experiments starting from *in vivo* or *in vitro* oral phase and ends in *in vitro* gastro-intestinal phase. Each digestion experiment was fed with one of five test substrates: As controls, cookie (a protein-free blank), Mix and Mix+maca. The three formulae: Ref+ref, Maca+ref and Maca+OLE. Two sets of experiments were conducted for each product sample: the first set included three replicate experiments with oral contributions from a single individual, while the second set involved three replicate experiments with contributions from three different individuals. SSF, simulated salivary fluid, SGF, simulated gastric fluid, SIF, simulated intestinal fluid. Created with BioRender.com

Table 2. Composition of stock solutions and simulated digestive fluids. Simulated salivary fluid (SSF), simulated gastric fluid (SGF), Simulated Intestinal fluid (SIF) used in *in vitro* adult and senior digestion experiments.

Compound	MW (gr/mol)	Stock [g/l]	[M] in stock	Volumes to add from stock solutions to generate 1 (L) solution					
				SSF [mL]*		SGF [ml]*		SIF [ml]*	
				Adult	Senior	Adult	Senior	Adult	Senior
KCl	74.55	46.72	0.627	24.1	10	11	56	10.8	10.8
KH ₂ PO ₄	136.09	68	0.5	7.4	40	1.8	1.8	1.6	1.6
NaHCO ₃	84.01	84	1	13.6	8	25	13	85	85
NaCl	58.44	120	2.053	-	2	23.6	20	19.2	16
MgCl ₂ (H ₂ O) ₆	203.3	30	0.148	1	2	0.8	4	2.2	2.2
(NH ₄) ₂ CO ₃	96.086	27.28	0.283	0.21	-	1.8	2	-	-
Urea	60.06	22.5			10		0.6		4.8

2.5.1 Sodium – Dodecyl Sulfate-Polyacrylamide Gel Electrophoresis (SDS-PAGE)

Differences in proteins digestion and peptides formation under the various digestion conditions were based on SDS-PAGE. This qualitative analysis was performed to enable determining protein dissipation during digestion alongside monitoring the breakdown patterns formed therein. The SDS-PAGE analysis followed the method described previously[74] with slight modifications. Samples were loaded onto 12%

acrylamide gels and focused on the molecular weight range between 10-170 kDa using the protein size standard Spectra Multicolor Low Range Protein Ladder (Thermo Scientific).

2.5.2 Proteomic analysis of bioaccessible peptides

Liquid chromatography-tandem mass spectrometry (LC-MS/MS) analyses were used to identify the bioaccessible peptides found in aspirated digesta samples. The protein samples were first subjected to 8.5M Urea, 100mmol/L ammonium bicarbonate, and 10mmol/L DTT, followed by vortexing and centrifugation at 16,000g for 15 minutes at room temperature. Protein amount estimation was conducted using Bradford readings.

The samples underwent reduction (60°C for 30 min) and modification with 35.2mmol/L iodoacetamide in 100mmol/L ammonium bicarbonate (room temperature for 30 min in the dark). Post-modification, samples were diluted to 6M Urea and filtered using Omicon 10kD (centrifugal filter Amicon® Ultra, 0.5 mL, 10 kDa). Peptides from the filtrate were desalted using a homemade C18 stage tip, dried, and resuspended in 0.1% Formic acid, 2% acetonitrile. Peptides were analyzed by LC-MS/MS using a Q Exactive HF mass spectrometer (Thermo) coupled with a capillary HPLC (Evosep). The peptides were loaded onto an 8cm, ID 150µm, 1.9-micron Endurance column EV1109 (Evosep) and eluted with the built-in Xcalibur 30 SPD method (44 min). Mass spectrometry was performed in positive mode, involving repetitively full MS scans (m/z 300–1500) and High-energy Collision Dissociation (HCD) of the 18 most dominant ions (>1 charges) selected from the full MS scan. Dynamic exclusion with a 20s duration was enabled. Protein Discoverer 2.4 (Thermo) with the Sequest search engine was utilized for mass spectrometry data analysis. The search encompassed proteomes from various sources, including UniProt and NCBI nr databases, with specified mass tolerances. Oxidation on methionine and protein N-terminus acetylation were considered as variable modifications, while carbamidomethyl on cysteine was considered a static modification. Label-free analysis was employed for data quantification, and peptide-level false discovery rates (FDRs) were filtered to 1% using the target-decoy strategy amino acids).

All samples were found to contain over 1200 different peptides sequences. The peptides sequences were classified according to their length and the percentage of each length group was evaluated in each sample.

2.5.3 *In vitro* digestible indispensable amino acid score (DIAAS)

In vitro digestibility and DIAAS determinations were carried out as described previously[77]. Similar experiments were also conducted on samples subjected to *in vitro* senior digestion to evaluate total *in vitro* digestibility and DIAAS values under these conditions[9]. A protein-free blank (cookie) was digested in parallel to serve as control and all samples were subjected to analyses in triplicate. The total amino acids (AAs) of each substrate were determined as described in ISO 13903:2005 [78]. Samples were hydrolyzed for 24h with 6M HCl, then derivatized with AccQ-Tag Ultra reagent (Waters, 2007) before the amino acid (AA) profile was determined by ultra-high-performance liquid chromatography (UHPLC) (AccQ-Tag Ultra 2.1 × 100 mm, 1.7 µm, Waters) coupled with a UV detector (Vanquish, Thermo Scientific, Reinach, Switzerland). Digestibility percentages for both total and individual AAs of the substrates subjected to IVD were determined

through measuring the digestible fraction of the food (supernatant) and dividing it by the total food aliquot (supernatant + pellet [undigested fraction, according to the formula (1):

$$(1) \text{ In vitro digestibility } [\%] = \frac{Fs - Cs}{(Fs - Cs) + \max(0; Fp - Cp)} * 100$$

Fs = Food supernatant, Cs = Cookie supernatant, Fp = Food pellet, Cp = Cookie pellet.

Then, the calculation of *in vitro* digestible indispensable amino acid (*in vitro* DIAA) per gram of food protein was calculated according to the formula (2):

$$(2) \text{ in vitro DIAA} = \text{mg of IAA per g of food protein} \times \text{in vitro digestibility of IAA}$$

The *in vitro* DIAA ratio (DIAAR) per gram of food protein was calculated, considering the reference protein for older child, adolescent, adult, given by FAO [79], according to the formula (3):

$$(3) \text{ in vitro DIAAR } (\%) = \frac{\text{mg of in vitro DIAA in 1 g of dietary protein}}{\text{mg of the same dietary IAA in 1 g of the reference protein}}$$

DIAAR was computed for each individual indispensable amino acid (IAA) and the DIAAS for a specific food corresponds to the lowest DIAAR among them.

2.6. Statistical analysis

All experiments were conducted in duplicates or triplicates, and the results are presented as the calculated mean and standard deviation. Statistical analyses were performed using various methods tailored to specific measurements. For general statistical analyses, a one-way ANOVA ($P < 0.05$) with a Tukey's Multiple Comparison test and Kruskal Wallis H Test were carried out using MATLAB and Excel. For Tribological measurements and Microstructure characterization of products, a one-way ANOVA followed by the Tukey's pairwise comparison with a 95% confidence interval was performed using SPSS software (IBM, SPSS Statistics, version 24).

The TDS was illustrated as dominance curves vs. time, employing standardized times (from T_0 to T_{100}). Two crucial levels were outlined for interpretation: the "chance level" (P_0), representing the dominance rate an attribute could obtain by chance, and the "significance level" (P_s), denoting the minimum dominance rate for the attribute occurrence to be considered significantly higher than the chance level P_0 [80]. For the analysis of QDA data, an ANOVA with assessor (random) and product (fixed) effects was applied. The Tukey post hoc test was utilized to compare attribute means among different products.

3. Results and discussion

In line with the goal of the EAT4AGE project, this work targeted the development of a novel food product that is rationally designed to tackle the nutritional challenges of old age. In essence, the work explored the production and characteristics of the products, the perception of their palatability in the target population and their potential digestive fate in seniors. Following product design, formulation and optimization of the extrusion conditions, three plant-based cereals were manufactured and tested in the prospect of healthy community-dwelling seniors.

3.1. In-vitro characterization of the plant-based cereal prototypes

3.1.1. Proximate analyses

This work developed food prototypes that could be classified as ultraprocessed foods whose association with adverse effects to health are vividly discussed these days [81], [82], [83], [84]. Yet, proximate analyses summarized in **Table 3** confirmed that the prototypes boast a protein content exceeding 12% (w/w), a fat content of approximately 20%, and minimal sugar (less than 5%). Interestingly, this nutritional profile surpasses that of commercially available counterparts, which typically offer around 6% (w/w) protein, ~15% fat and a very high sugar content of approx. 40% (w/w). Accumulating evidence on the beneficial role of dietary fiber and whole grains [85], [86], [87], [88], [89], [90] was also addressed in the product formulation which was calculated to contain 17% (w/w) dietary fiber. Further, studies highlight the important role of micronutrients like Iron (Fe), Zinc (Zn), Copper (Cu), and Selenium (Se) in the nutrition of older adults [5], [15], [91]. Nevertheless, cereals often fall short in providing adequate quantities of these essential nutrients. Thus, the inclusion of legumes, like chickpeas, offered an opportunity to compensate for the AA profile and this inherent drawback of micronutrient content of cereals, namely as it pertains iron content as depicted in **Table 3**. Combining these complementary sources enhances the overall nutritional profile of the extruded cereal products, thus, highlighting the importance of product formulation over processing [81]. However, the success of such novel and healthier food choices also mandates matching their nutritional potential with consumer adoption that stems from the palatability.

3.1.2. Instrumental single compression test

One of the major sensorial traits of cereal products is their distinct texture and descriptors, such as crunchiness, chewiness and sogginess which can be elemental in the development of future foods for seniors [10]. As a first step, instrumental single compression measurements were initiated and sought to measure the impact of supplementation on the mechanical properties of the products (**Figure 2**). These analyses revealed that addition of the functional supplements (Maca and OLE) significantly lowered sample hardness ($P < 0.05$) to values lower than that reported for an extrudate of maize-millet fortified with soy flour [92]. This would imply a possible improvement in their chewiness and oral comfort for seniors. In addition, the supplementation of Maca involved a reduction in corn grits content (**Table 1**) and product hardness (**Figure 2**) which coincides with recent studies correlating diminished grits content with a subsequent reduction in product hardness [93], [94]. These findings underscore the intricate influence of formulation adjustments on the characteristic features of snacks. Furthermore, the examination reveals minimal variance in young's modulus among the tested products (**Figure 2B**). In comparison to other items, our product demonstrates a lower stiffness than snacks [95] and crackerbread [96], again implying a propensity for improved oral comfort in older adults. Thus, further experiments were held to improve the insight into the oral processing and palatability of the products.




Figure 2. Texture Profile Analysis (TPA) for plant-based prototypes- Enhancing oral comfort through functional materials Addition. (A) Strength to strain TPA graph (B) Summary table of textural properties values. Values (mean ± SD) in the same line followed by a different letter are significantly different ($P < 0.05$).

3.1.3. Characterization of samples through tribological measurements

The lubrication properties of the products were evaluated using a ball-on-disk tribometer featuring hydrophobic elastomer surfaces resembling bio-tribological contacts (**Figure 3**). The Stribeck curve graphically illustrates the tribological behavior and lubrication properties, traditionally divided into three regimes: boundary, mixed, and hydrodynamic.

The friction behavior of water shows that lubrication occurred in the boundary and mixed regimes, whereas the lubrication behavior of model saliva showed boundary, mixed and the onset of hydrodynamic regime, indicating that both lubricants and the surface properties influenced the lubrication properties. For Maca+OLE boli, the friction curve displayed the three regimes, a boundary plateau at lower speeds (<10 mm/s), a mixed regime (10 – 100 mm/s), and a hydrodynamic lubrication regime (>100 mm/s) (**Figure 3**). In contrast, both Ref+ref and Maca+ref boli exhibited superimposed behavior with low μ at entrainment speeds <100 mm/s, corresponding to the mixed regime, and an increase in μ when entrainment speeds exceeded 100 mm/s, indicative of the hydrodynamic lubrication regime (**Figure 3**). These frictional curve behaviors mirror the lubrication characteristics observed in some edible oils and plant-derived oleosomes [97], [98] suggesting

that some oil might have been released from the cereal matrix. Considering that all products had similar composition ($P > 0.05$) (Table 3), the differences in oil release and its effect on the lubrication profile might have been influenced by the fracture properties of the products, where Maca+OLE presented significantly reduced sample hardness ($P < 0.05$) (Figure 2). The impact of mechanical properties on chewing and bolus formation has been previously discussed, noting that the formation of larger bolus particles tends to decrease as the hardness of food increases [99]. Consequently, it is plausible that a reduced breakdown of the food matrix could contribute to a reduction in the release of oil from the matrix.




Figure 3. Mean friction coefficients as a function of entrainment speeds and CLSM images of (A) Ref+ref, (B) Maca+ref, (C) Maca+OLE. Protein aggregates are stained fluorescently green using Fast green whilst fat droplets are stained fluorescently red using Nile Red. The friction curve of water and model saliva are added as reference. Data represent average of triplicate measurements on three samples ($n = 2 \times 3$).

Table 3. Proximate analyses of plant-based cereal prototypes ($n=3$) (30 g portion size)

	Pre-extrusion		Post-extrusion		
	Mix	Mix+maca	Ref+ref	Maca+ref	Maca+OLE
Calories* (kcal)			~126	~126	~126
Total fat (gr)			5.112 ± 0.006 ^a	5.7 ± 0.2 ^a	6.0 ± 0.3 ^a
Protein (gr)	3.15 ± 0.06 ^a	3.24 ± 0.03 ^a	3.83 ± 0.02 ^b	3.75 ± 0.06 ^b	3.9 ± 0.1 ^b
Sodium (gr)	0.03 ± 0.02	0.08 ± 0.03	0.06 ± 0.03	0.06 ± 0.02	0.06 ± 0.02
Total Carbohydrate* (gr)			8	8	8
Dietary fiber* (gr)			5 (20%)	5 (20%)	5 (20%)
Total sugar (gr)			1.3 ± 0.2 ^a	1.0 ± 0.2 ^a	1.2 ± 0.2 ^a
Iron (mg)	9 ± 2	8 ± 1	6.8 ± 1.8 (85%)	5.0 ± 0.2 (63%)	4.0 ± 0.3 (50%)
Calcium (mg)	32.99 ± 2.91	51 ± 7	32 ± 5 (3%)	37.90 ± 2.65 (3%)	32 ± 2 (2%)
Magnesium (mg)	30.2 ± 3.1	41 ± 7	40 ± 5 (11%)	49 ± 4 (13%)	41 ± 4 (11%)
Phosphorus (mg)	36 ± 3	49 ± 5	60 ± 4 (9%)	78 ± 3 (11%)	67 ± 4 (10%)
Potassium (mg)	93.20 ± 9.22	150 ± 18	108.76 ± 11.78 (4%)	142.50 ± 6.81 (5%)	108.3 ± 9.3 (4%)

Zinc (mg)	0.66 ± 0.07	0.59 ± 0.05	0.94 ± 0.31 (10%)	0.9650 ± 0.2951 (10%)	0.9 ± 0.3 (10%)
Copper (mg)	0.13 ± 0.05	0.092 ± 0.008	0.3 ± 0.1 (0%)	0.164 ± 0.011 (0%)	0.14 ± 0.02 (0%)
Selenium (mg)	19 ± 2	24.937 ± 4.281	23.71 ± 5.31 (43%)	32 ± 4 (58%)	28 ± 4 (51%)
Manganese (mg)	1.0 ± 0.6	0.53 ± 0.05	0.9173 ± 0.4621 (45%)	0.48 ± 0.02 (24%)	0.31 ± 0.05 (15%)

Values (mean ± SD) in the same line followed by a different letter are significantly different ($P < 0.05$). *Theoretical values based on individual ingredients composition (“Tzameret” software program used by the Israeli Ministry of Health). **(**Value%**)-Percentage of average RDA for men and women aged 51-70 ***Mix, Mix+maca are the pre-extruded products with the same formulation of Ref+ref and Maca+OLE respectively.

As determining bolus particle size using the Mastersizer instrument was considered inappropriate due to potential shear degradation caused by the impeller, and because conventional laser diffraction instruments assume spherical particles leading to an overestimation, especially for the volume-weighted mean diameter, which emphasizes larger particles within the distribution, confocal microscopy images of the bolus samples were acquired to observe the relative distribution of protein and fat components (**Figure 3**). In both, Ref+ref and Maca+ref boluses, substantial protein structure fragments and prominent fat coalescence areas were evident. In contrast, Maca+OLE bolus displayed smaller protein fragments and more uniformly distributed smaller oil droplets. Based on these observations, it is likely that in both Ref+ref and Maca+ref boluses, the observed low μ at entrainment speeds <100 mm/s is associated with the formation of a lubricating oil layer on the surfaces. At entrainment speeds exceeding 100 mm/s, some small particles might have entrained into the contact, becoming dominant and resulting in higher μ . For Maca+OLE the friction behavior was likely governed by the small, dispersed oil droplets and soluble substances forming the boundary lubricating regime. As speed increased (>10 mm/s), bigger boli particles/ droplets might entrained between contact surfaces, forming a lubricating film and decreasing μ with increasing speed. At high entrainment velocities (>100 mm/s), complete separation of surfaces occurred, and large particles entering the gap increased the friction coefficient (**Figure 3**). Thus, these experiments helped quantitate some of the characteristics of oral boli of the products and establish the possible benefits that arise from the use of the functional supplements. Yet, these instrumental measures needed to be confirmed in the target population through corresponding sensory evaluations of oral comfortability and palatability.

3.2. Comprehensive in vivo sensory evaluation

3.2.1. Food comfortability

Mastication is a key step in the indulgence humans extract out of food and it has repercussions to the product’s commercial success as well as its subsequent breakdown in the consumer’s gut. Thus, food comfortability was evaluated in a human cohort using a questionnaire following a procedure described previously [68], [100] and the results are plotted on different spider plots (**Figure 4**). First, self-reports on bolus formation (**Figure 4A**), mouth pain (**Figure 4B**), texture sensation (**Figure 4C**) and flavor (**Figure 4D**) were assessed and found not to be adverse nor modulated significantly by the addition of the supplements. Particularly noteworthy is the observed self-reported ease in performing important masticatory operations (cutting, chewing, and swallowing) towards bolus formation, coupled with a relatively short total time (**Figure 4A**) and very low reports on mouth pain (**Figure 4B**). Together with the instrumental texture results (**Figure 2**), the findings indicate enhanced oral comfort of the functional product which could be particularly beneficial

for seniors with physical limitations hindering the consumption of complex or tough foods[5]. Analysis of the perceived properties of the products reveal that the products were characterized as being crunchy and melting (Figure 4C). This suggests that the filling does not compromise the shell integrity and hardness due to possible moistening. The absence of extreme flavors (Figure 4D) and low scores for bitterness and acidity, coupled with the low sugar content (Table 3), may contribute to this outcome. These positive perceived properties were also expressed in the high scoring of the products' oral comfort (Figure 4E), with scores exceeding 4.8 out of 5.0. This adds an extra layer of evidence to reinforce the overall positive reception of the products and their potential palatability to the target customers.




Figure 4. Results from the questionnaire on the perception of food comfortability. (A) Bolus Formation and Time Needed (Scale: "Impossible" to "Very Easy" and "Impossible" to "Very Brief"), **(B)** Pain Perception (Scale: "Extremely" to "Not at All"), **(C)** Texture Perception (Scale: "Extremely" to "Not at All"), **(D)** Flavor Perception (Scale: "Extremely" to "Not at All"), **(E)** General Food Comfort (Scale: "Very Uncomfortable" to "Very Comfortable"). Values followed by a different letter are significantly different ($P < 0.05$). All sections consistently use a 5-point scale.

3.2.2. Sensory descriptions by the trained panel via Quantitative Descriptive Analysis (QDA) and Temporal dominance of sensations (TDS)

QDA provided a comprehensive description of sensory properties from a static perspective, while TDS identified the dominant properties over the consumption period (dynamic perception). The outcomes demonstrated were very much aligned, with minor differences observed among the samples (**Figure 5**).




Figure 5. TDS, QDA, and Spider plots derived from the trained panel sensory description of two products (Maca+ref, Maca+OLE). (A) TDS curve of Maca+ref, (B) TDS curve of Maca+OLE, (C) QDA results of the 2 formulae (Values followed by a different letter are significantly different ($P < 0.05$)), (D-F) Spider plots of QDA results (*Significant $P < 0.05$ **Significant $P < 0.01$ ***Significant $P < 0.001$).

According to TDS (**Figure 5A-B**), crispiness — a desirable characteristic for this product- dominated at the beginning of oral processing for Maca+ref and Maca+OLE, persisting for approximately 50% of the eating duration. Chocolate flavour prevailed in the middle of the consumption process for all samples, especially Maca+OLE (**Figure 5C**). Toward the end, bitterness and stickiness became more prominent, although these attributes received low to medium scores (0.2–0.4), suggesting either limited consensus among evaluators, or that there were no clear dominating attributes in the evaluation. The results indicated that the combination of flours and the oil-based filling resulted in a somewhat sticky bolus, occurring toward the end of consumption. However, this was not a major concern, as research suggests that boluses with more saliva tend to have a dry and sticky texture, which can be beneficial for oral issues in the older adult demographic [101]. Moreover, prolonged chewing associated with sticky foods is essential for both eating and overall well-being. Recent research highlights the cognitive benefits of chewing, particularly in the hippocampus, a region of the central

nervous system responsible for spatial memory and learning [102]. The panel described the Maca sample as significantly less roasted, chocolatey and less juicy, and with significantly more drawer odour (**Figure 5**). The older adult panel, however, did not notice the differences in dryness (**Figure 4**), which makes sense as the trained panel is more sensitive because of the training, and they found a significant but minor difference. In terms of bitterness, Maca and OLE are usually bitter and earthy flavours, the panel rated both samples similarly in terms of bitterness. However, their inclusion had no discernible effect on product acceptance or characteristics, as supported by the tasting tests conducted on older adult participants in Israel (**Figure 4**). The trained panel evaluated the samples in accordance with the data of oral comfort as described by the older adult subjects, revealing minimal significant differences among various samples.

3.3. Studying protein breakdown using a semi-dynamic in vitro digestion model

3.3.1. Sodium – Dodecyl Sulfate-Polyacrylamide Gel Electrophoresis (SDS-PAGE)

Age-related changes have been shown to affect digestive parameters [5]. In turn, altered GI functions may divert the digestive fate of foods compared to that assessed in healthy young adults. Therefore, this work evaluated the IVD proteolysis of the products using a recently agreed protocol for recreating digestion in older adults [9]. Utilizing SDS-PAGE analysis as a primal qualitative tool, we dissected protein digestion of the various products and compared between their digestion in young and older adults (**Figure 6** and **Figure 7**, respectively). First, we investigated the impact of adding functional ingredients (Maca + OLE) on protein breakdown trajectories in seniors. Studies indicate that food antioxidants may interact with proteins thereby hindering their accessibility to digestive breakdown [47], [48]. The findings here (**Figure 6**) revealed no significant differences in the protein breakdown patterns, with a sustained gastric digestion that is practically completed during the intestinal phase. Moreover, bands corresponding to most proteins and large peptides, observed during the gastric phase, had dissipated during the intestinal phase. Yet, extrusion processing may impact protein nutritional quality, with studies highlighting both beneficial and deleterious effects of processing on protein digestibility [37], [50]. The structural changes induced by high-temperature, short-duration, and high-pressure conditions of extrusion may enhance protein digestibility by denaturing proteins and exposing enzyme-accessible sites or inactivation of anti-nutritional moieties [50]. Thus, further tests focused on the protein digestibility of pre- and post-extruded plant-based cereals, as determined from analysis of the bioaccessible soluble aliquots of digestive effluents (**Figure 7**). It is important to note, the pre-extruded product (Mix+maca) exhibited faint protein bands that were attributed to the lack of release of the proteins from the food matrices, i.e. proteins were not released to the gastric lumen or collected for the SDS-PAGE analyses, both in young and older adults (**Figure 7A** and **7C**). Contrary, visible protein bands were noted in the corresponding gels of processed samples (**Figure 7B** and **7D**), which suggests a higher bioaccessibility of proteins in these processed samples, both for young and older adults. This would concur with previous studies supporting extrusion processing potential to unleash the nutritional potential of plant proteins [36], [37], [50]. When comparing the digestive trajectories observed during digestion of the processed samples (**Figure 7B** and **7D**), protein breakdown was found to be more delayed under the senior's gut

conditions compared to that of a young adult. In fact, protein bands were found to endure no longer than 30 min of gastric digestion under the conditions of a young adult (**Figure 7D**) while such bands were found to persist for up to 180min of gastric digestion under conditions of older adults (**Figure 7B**). This noticeable divergence between the two target populations emerged within just 15 minutes of gastric digestion, signifying a delayed digestion in seniors. This distinction underscores the influence of physiological changes on the digestive proteolysis patterns of food products in young and older adults which could have nutritional ramifications. Therefore, further in-depth qualitative view of protein breakdown profiles were sought.




Figure 6. Consistent protein breakdown pattern: SDS-PAGE analyses of senior digestion for post-extruded products. (A) Ref+ref, (B) Maca+ref, (C) Maca+OLE.




Figure 7. Comparative SDS-PAGE analyses of digesta samples: Enhanced proteolysis in post-extruded products (Maca+OLE) and reduced proteolysis in senior *in vitro* digestion vs. adult. (A+C) Protein breakdown patterns of the pre-extruded (Mix+maca) (B+D) and post-extruded product (Maca+OLE) under senior (A+B) and adult (C+D) *in vitro* digestion conditions.

3.3.2. Proteomic analysis of bioaccessible peptides

From the nutritional perspective, proteins are broken down into bioaccessible peptides and AAs that are up taken into the body where they may exercise their biological function. First, LC-MS/MS analysis was used to identify the bioaccessible peptides present in aspirated digesta samples (**Figure 8**). These analyses provide a comparative look into the bioaccessible peptide distributions recovered from the digestion of pre-extrusion and post-extrusion products as well as differentiating between the adult and older adults' patterns, according to peptide AA length. These revealed no significant differential patterns of bioaccessible peptides between adults and seniors (**Figure 8**). This could be attributed to the dense matrix of cereal products that may limit pepsin's access to protein substrates, as reported in the past for proteinaceous gels [103], [104]. The intricate nature of the cereal product matrix may obscure physiological variations between adults and the older adults, potentially making any differences in digestibility less conspicuous in our findings. While SDS-PAGE analyses highlighted a delayed proteolysis in seniors for proteins between 170 kDa-10kDa, these analyses indicate that at the low MW peptide level below 10 kDa (up to 52AA) there are no pronounced differences.

A similar investigation comparing meat digestion in seniors and adults yielded analogous results [43], supporting the notion that the properties of the food matrix may obscure effects arising from age-related physiological differences in gut functions. However, two points should be emphasized. One, the absence of apparent differences noted herein do not necessarily imply equal digestion, as the IVD findings do not account for the full kinetics of digestive breakdown [105], [106]. Second, similarities in the bioaccessible peptide lengths does not necessarily mean similarities in the AA sequences of these peptides. Moreover, these analyses focused on peptide lengths between 10AAs up to 52AAs, thus, overlooking insights into free amino acids (FAAs) which are absorbable and of nutritional importance [107], [108], [109], [110].




Figure 8. Uniform peptide distribution across LC-MS/MS samples collected from (A) The end of the gastric phase and (B) The end of the intestinal phase in IVD model under senior and adult conditions of the different formulae. Columns for adult are in full color, and for senior are dashed.

3.3.3. *In vitro* digestible indispensable amino acid score (DIAAS)

From the nutritional perspective, protein quality is assessed through scoring its ability to provide the body with amino acids. Practically, this is done through various methods with DIAAS being the most recently advocated method [77], [79], [111], [112]. The newly introduced *in vitro* DIAAS methodology provides a robust framework for assessing the protein quality of human foods, leveraging ileal digestibility as a key metric and circumventing the need for ethically challenging and time-consuming human trials [77]. This *in vitro* scheme was used in this study and the resulting digestibility data of the functional product (before and after processing) are given in **Figure 9 and supplementary Table 2S**. Comparing the product's digestibility in adults or seniors (over 65 years) gave rise to two main observations. One was that there were no significant differences in the product's digestibility and nutritional quality in adults or seniors (**Figure 9A and 9B**). The second observation is that processing increased the product's digestibility to obtain highly digestible proteins (total protein digestibility exceeding 80%, **Figure 9B**). This showcases the potential for producing a nutritionally rich cereal product with sustainable plant-based protein sources. Moreover, these digestibility values surpass the typical digestibility range of plant-based sources, which are typically between 40% and 80% [36], [37], [77].




Figure 9. High *In vitro* Digestibility highlighted in results. (A) Comparative protein digestibility percentages in different cereal products pre- and post-extrusion under adult and senior IVD models, (B) Table presenting digestibility percentages of individual AAs. Values are expressed as the mean \pm SD. Different letters denote significant differences ($P < 0.05$)

Furthermore, the DIAAS values were determined by considering the lowest *in vitro* DIAA ratios for older child, adolescents, and adults for each analyzed substrate. Computed using the formulae (2+3), the resulting lowest limiting AA and their respective value were as follows: In the pre-extruded product, Mix+maca exhibited *in vitro* DIAAS values of Lysine (84 ± 46) and Tryptophan (64 ± 35) for adults and seniors, respectively. For the post-extrusion product, Maca+OLE recorded values of Lysine (34 ± 3) and

Lysine (33.9 ± 0.3) for adults and seniors, respectively. These findings highlight that co-extrusion may decrease the bioaccessibility of specific AAs, notably lysine and tryptophan which concurs with previous reports [50], [113]. This affirmed that although processing may compromise the bioaccessibility levels of specific AAs, this did not come at the expense of an overall low digestibility of the product. Thus, this work confirmed that the combined formulation of a cereal product with four different flours yielded a palatable product with high digestibility scores, even when assessed under the age- deteriorated conditions of the older adults' gut.

4. Conclusion

Nutrition is expected to play a pivotal role in promoting healthy aging, and age-tailored functional foods could help capitalize on this notion. To this end, food research and development efforts need to support dietary shifts towards healthier and more sustainable food choices, such as those suggested in the EAT-Lancet report [4]. As an example of relevant EAT4AGE consortium outputs, this work developed plant-based prototypes of co-extruded cereals designed for superior macronutrient profiles that surpass existing commercial products and are pleasurable to the consumer. With *in vitro* digestibility exceeding 80%, this project makes a valuable addition to the table of choices for a balanced and sustainable diet for the aging demographic. Yet, bridging the gap between our various analyses and real-world *in vivo* efficacy requires further investigation which could be jointly pursued by industry and academia together. In fact, this work joins rising efforts to develop nutritional solutions that seek to tackle concerns over the health gap of the soaring global aging population. Overall, this research emphasizes a nutritional paradigm that tailoring food solutions for seniors, or any specific target population, need to intersect nutritional values with consumer preferences for enjoyment. Thus, this work hopes to stimulate an avenue of endeavors towards innovative, nutritionally optimized products that seek to boost human health and quality of life.

Ethical statement

The Technion Ethics Committee granted this research its institutional review board (IRB) permission (form No. 139-2022) and approved the untrained consumer panel as well as the ethical collection of human salivary and bolus samples as part of it (both conducted in Israel). Each participant signed informed permission papers after being made aware of the study's objectives and methods, being checked to see if they met the requirements for inclusion and passing a questionnaire.

Declaration of competing interest

The authors report no conflict of interest.

Acknowledgments

We acknowledged the financial support of the Israeli Ministry of Science and Technology Grant #3-16518 and the Research Council of Norway (project 321860), under the umbrella of the European Joint Programming Initiative "A Healthy Diet for a Healthy Life" (JPI HDHL) and of the ERA-NET Cofund

ERA-HDHL (GA N° 696295 of the EU Horizon 2020 Research and Innovation Programme)”; as well as the support of the Russell Berrie Nanotechnology Institute (Technion – Israel Institute of Technology, Haifa, Israel). Ms. L. Mashiah would also like to thank the partial support of her PhD by the Israel Scholarship Education Foundation (ISEF). Special thanks are also extended to the staff of the Smoler Proteomics Center at the Technion for their support and collaboration. Authors P.Varela and J.Skaret also wish to thank the Norwegian Fund for Research Fees for Agricultural Products (FFL) for support, through the project “FoodForFuture” (Project number 314318; 2021-2024). A.Sarkar and A. Araiza Calahorra acknowledge the EAT4AGE project fundings received from [Biotechnology and Biological Sciences Research Council (BBSRC), UK: BB/V018914/1] under the umbrella of the European Joint Programming Initiative “A Healthy Diet for a Healthy Life” (JPI HDHL) and of the ERA-NET Cofund ERA-HDHL (GA No 696295 of the EU Horizon 2020 Research and Innovation Programme).

References

- [1] W. He, D. Goodkind, and P. Kowal, ‘An Aging World: 2015’.
- [2] C. E. Brett, D. Dykiert, J. M. Starr, and I. J. Deary, ‘Predicting change in quality of life from age 79 to 90 in the Lothian Birth Cohort 1921’, *Quality of Life Research*, vol. 28, no. 3, pp. 737–749, Mar. 2019, doi: 10.1007/S11136-018-2056-4/TABLES/4.
- [3] A. Green, C. Blattmann, C. Chen, and A. Mathys, ‘The role of alternative proteins and future foods in sustainable and contextually-adapted flexitarian diets’, *Trends Food Sci Technol*, vol. 124, pp. 250–258, Jun. 2022, doi: 10.1016/J.TIFS.2022.03.026.
- [4] W. Willett *et al.*, ‘Food in the Anthropocene: the EAT–Lancet Commission on healthy diets from sustainable food systems’, Feb. 02, 2019, *Lancet Publishing Group*. doi: 10.1016/S0140-6736(18)31788-4.
- [5] D. Rémond *et al.*, ‘Understanding the gastrointestinal tract of the elderly to develop dietary solutions that prevent malnutrition’, *Oncotarget*, vol. 6, no. 16, p. 13858, Jun. 2015, doi: 10.18632/ONCOTARGET.4030.
- [6] C. L. Bush *et al.*, ‘Toward the Definition of Personalized Nutrition: A Proposal by The American Nutrition Association’, 2019, doi: 10.1080/07315724.2019.1685332.
- [7] E. Biagi, C. Franceschi, S. Rampelli, M. Capri, P. Brigidi, and M. Candela Correspondence, ‘Gut Microbiota and Extreme Longevity’, *Current Biology*, vol. 26, pp. 1480–1485, 2016, doi: 10.1016/j.cub.2016.04.016.
- [8] M. J. Claesson *et al.*, ‘Gut microbiota composition correlates with diet and health in the elderly’, *Nature* 2012 488:7410, vol. 488, no. 7410, pp. 178–184, Jul. 2012, doi: 10.1038/nature11319.
- [9] O. Menard *et al.*, ‘Static in vitro digestion model adapted to the general older adult population: an INFOGEST international consensus’, *Food Funct*, 2023, doi: 10.1039/D3FO00535F.
- [10] A. Sarkar, ‘Oral processing in elderly: understanding eating capability to drive future food texture modifications’, *Proceedings of the Nutrition Society*, vol. 78, no. 3, pp. 329–339, Aug. 2019, doi: 10.1017/S0029665118002768.
- [11] P. T. Morgan, O. C. Witard, G. Højfeldt, D. D. Church, L. Breen, and P. Morgan, ‘Dietary protein recommendations to support healthy muscle ageing in the 21st century and beyond: considerations and future directions’, *Proceedings of the Nutrition Society*, pp. 1–14, doi: 10.1017/S0029665123003750.
- [12] F. B. Hu, ‘Key Symposium Diet strategies for promoting healthy aging and longevity: An epidemiological perspective’, 2023, doi: 10.1111/joim.13728.
- [13] E. Dent, O. R. L. Wright, J. Woo, and E. O. Hoogendijk, ‘Malnutrition in older adults’, *The Lancet*, vol. 401, no. 10380, pp. 951–966, Mar. 2023, doi: 10.1016/S0140-6736(22)02612-5.
- [14] J. A. Phillips, ‘Dietary Guidelines for Americans, 2020–2025’, <https://doi.org/10.1177/21650799211026980>, vol. 69, no. 8, p. 395, Jul. 2021, doi: 10.1177/21650799211026980.
- [15] M. J. Bruins, P. Van Dael, and M. Eggersdorfer, ‘The role of nutrients in reducing the risk for noncommunicable diseases during aging’, Jan. 01, 2019, *MDPI AG*. doi: 10.3390/nu11010085.

- [16] C. Putra, N. Konow, M. Gage, C. G. York, and K. M. Mangano, 'Protein Source and Muscle Health in Older Adults: A Literature Review', *Nutrients* 2021, Vol. 13, Page 743, vol. 13, no. 3, p. 743, Feb. 2021, doi: 10.3390/NU13030743.
- [17] F. B. Hu, 'Diet strategies for promoting healthy aging and longevity: An epidemiological perspective', *J Intern Med*, vol. 295, no. 4, pp. 508–531, Apr. 2024, doi: 10.1111/JOIM.13728.
- [18] A. K. Gosby *et al.*, 'Testing protein leverage in lean humans: A randomised controlled experimental study', *PLoS One*, vol. 6, no. 10, Oct. 2011, doi: 10.1371/journal.pone.0025929.
- [19] A. K. Gosby, A. D. Conigrave, D. Raubenheimer, and S. J. Simpson, 'Protein leverage and energy intake', *Obesity Reviews*, vol. 15, no. 3, pp. 183–191, Mar. 2014, doi: 10.1111/OBR.12131.
- [20] E. A. Nunes *et al.*, 'Systematic review and meta-analysis of protein intake to support muscle mass and function in healthy adults', Apr. 01, 2022, *John Wiley and Sons Inc.* doi: 10.1002/jcsm.12922.
- [21] J. Aschemann-Witzel, R. F. Gantriis, P. Fraga, and F. J. A. Perez-Cueto, 'Plant-based food and protein trend from a business perspective: markets, consumers, and the challenges and opportunities in the future', <https://doi.org/10.1080/10408398.2020.1793730>, pp. 1–10, 2020, doi: 10.1080/10408398.2020.1793730.
- [22] R. Verkerk, 'EAT-Lancet-is there such a thing as "one-size-fits-all" sustainability?', 2019.
- [23] K. Gul, A. K. Singh, and R. Jabeen, 'Nutraceuticals and Functional Foods: The Foods for the Future World', *Crit Rev Food Sci Nutr*, vol. 56, no. 16, pp. 2617–2627, Dec. 2016, doi: 10.1080/10408398.2014.903384.
- [24] I. Siró, E. Kápolna, B. Kápolna, and A. Lugasi, 'Functional food. Product development, marketing and consumer acceptance—A review', *Appetite*, vol. 51, no. 3, pp. 456–467, Nov. 2008, doi: 10.1016/j.appet.2008.05.060.
- [25] C. M. Hasler, 'Functional foods: Their role in disease prevention and health promotion', *Food Technol*, vol. 52, no. 11, pp. 63–70, 1998.
- [26] S. Martín-Peláez *et al.*, 'Effect of olive oil phenolic compounds on the expression of blood pressure-related genes in healthy individuals', *Eur J Nutr*, vol. 56, no. 2, pp. 663–670, Mar. 2017, doi: 10.1007/s00394-015-1110-z.
- [27] M. S. Lee, H. W. Lee, S. You, and K. T. Ha, 'The use of maca (*Lepidium meyenii*) to improve semen quality: A systematic review', Oct. 01, 2016, *Elsevier Ireland Ltd.* doi: 10.1016/j.maturitas.2016.07.013.
- [28] J. Gan, Y. Feng, Z. He, X. Li, and H. Zhang, 'Correlations between Antioxidant Activity and Alkaloids and Phenols of Maca (*Lepidium meyenii*)', *J Food Qual*, vol. 2017, 2017, doi: 10.1155/2017/3185945.
- [29] Z. S. Hajdu, L. Lorántfy, N. Jedlinszki, K. Boros, J. Hohmann, and D. Csupor, 'QUALITY CONTROL OF MACA-CONTAINING (*LEPIDIUM MEYENII* WALP.) DIETARY SUPPLEMENTS', *Budapest Acta Alimentaria*, vol. 44, no. 3, pp. 461–467, 2015, doi: 10.1556/066.2015.44.0018.
- [30] H. O. Meissner *et al.*, 'Hormone-Balancing Effect of Pre-Gelatinized Organic Maca (*Lepidium peruvianum* Chacon): (III) Clinical responses of early-postmenopausal women to Maca in double blind, randomized, Placebo-controlled, crossover configuration, outpatient study', *Int J Biomed Sci*, vol. 2, no. 4, p. 375, Dec. 2006.
- [31] A. E. Clewell *et al.*, 'A Comprehensive Toxicological Safety Assessment of an Extract of *Olea Europaea* L. Leaves (Bonolive™)', <https://doi.org/10.1177/1091581815619764>, vol. 35, no. 2, pp. 208–221, Dec. 2015, doi: 10.1177/1091581815619764.
- [32] W. Sun *et al.*, 'Oleuropein improves mitochondrial function to attenuate oxidative stress by activating the Nrf2 pathway in the hypothalamic paraventricular nucleus of spontaneously hypertensive rats', *Neuropharmacology*, vol. 113, pp. 556–566, Feb. 2017, doi: 10.1016/J.NEUROPHARM.2016.11.010.
- [33] M. Cotacallapa-Sucapuca *et al.*, 'Extrusion process as an alternative to improve pulses products consumption. A review', *Foods*, vol. 10, no. 5, May 2021, doi: 10.3390/foods10051096.
- [34] A. Alcorta, A. Porta, A. Tárrega, M. D. Alvarez, and M. Pilar Vaquero, 'Foods for plant-based diets: Challenges and innovations', Feb. 01, 2021, *MDPI AG.* doi: 10.3390/foods10020293.
- [35] S. M. F. Bessada, J. C. M. Barreira, and M. B. P. P. Oliveira, 'Pulses and food security: Dietary protein, digestibility, bioactive and functional properties', *Trends Food Sci Technol*, vol. 93, pp. 53–68, Nov. 2019, doi: 10.1016/J.TIFS.2019.08.022.
- [36] A. G. A. Sá, Y. M. F. Moreno, and B. A. M. Carciofi, 'Food processing for the improvement of plant proteins digestibility', *Crit Rev Food Sci Nutr*, Nov. 2020, doi: 10.1080/10408398.2019.1688249.

- [37] V. Orlien, K. Aalaei, M. M. Poojary, D. S. Nielsen, L. Ahn , and J. R. Carrascal, 'Effect of processing on in vitro digestibility (IVPD) of food proteins', *Crit Rev Food Sci Nutr*, vol. 0, no. 0, pp. 1–50, Sep. 2021, doi: 10.1080/10408398.2021.1980763.
- [38] M. Gallego, J. M. Barat, R. Grau, and P. Talens, 'Compositional, structural design and nutritional aspects of texture-modified foods for the elderly', Jan. 01, 2022, *Elsevier Ltd.* doi: 10.1016/j.tifs.2021.12.008.
- [39] J. Makame, A. A. Nolden, and M. N. Emmambux, 'Texture properties of foods targeted for individuals with limited oral processing capabilities: the elderly, dysphagia, and head and neck cancer patients', Mar. 21, 2023, *Royal Society of Chemistry.* doi: 10.1039/d3fo00363a.
- [40] R. S. Gibson, 'The role of diet- and host-related factors in nutrient bioavailability and thus in nutrient-based dietary requirement estimates', *Food Nutr Bull*, vol. 28, no. 1 SUPPL. 1, 2007, doi: 10.1177/15648265070281S108.
- [41] A. Araiza-Calahorra, A. R. Mackie, G. Feron, and A. Sarkar, 'Can tribology be a tool to help tailor food for elderly population?', *Curr Opin Food Sci*, vol. 49, p. 100968, Feb. 2023, doi: 10.1016/J.COFS.2022.100968.
- [42] C. S. Levi and U. Lesmes, 'Bi-compartmental elderly or adult dynamic digestion models applied to interrogate protein digestibility', *Food Funct*, vol. 5, no. 10, pp. 2402–2409, Sep. 2014, doi: 10.1039/C4FO00478G.
- [43] S. Denis *et al.*, 'Digestion of cooked meat proteins is slightly affected by age as assessed using the dynamic gastrointestinal TIM model and mass spectrometry', *Food Funct*, vol. 7, no. 6, pp. 2682–2691, Jun. 2016, doi: 10.1039/C6FO00120C.
- [44] B. Walther *et al.*, 'REVIEW GutSelf: Interindividual Variability in the Processing of Dietary Compounds by the Human Gastrointestinal Tract', 2019, doi: 10.1002/mnfr.201900677.
- [45] E. Hern andez-Olivas, S. Mu oz-Pina, A. Andr s, and A. Heredia, 'Age-related gastrointestinal alterations of legumes and cereal grains digestibility Running title: Gastrointestinal alterations with the elderly of pulses and grains 2', 2021.
- [46] S. Lamothe, N. Azimy, L. Bazinet, C. Couillard, and M. Britten, 'Interaction of green tea polyphenols with dairy matrices in a simulated gastrointestinal environment', *Food Funct*, vol. 5, no. 10, pp. 2621–2631, Sep. 2014, doi: 10.1039/C4FO00203B.
- [47] T. Ozdal, E. Capanoglu, and F. Altay, 'A review on protein–phenolic interactions and associated changes', *Food Research International*, vol. 51, no. 2, pp. 954–970, May 2013, doi: 10.1016/J.FOODRES.2013.02.009.
- [48] J. Wang, S. Hu, S. Nie, Q. Yu, and M. Xie, 'Reviews on Mechanisms of in Vitro Antioxidant Activity of Polysaccharides', *Oxid Med Cell Longev*, vol. 2016, 2016, doi: 10.1155/2016/5692852.
- [49] A. Brodkorb *et al.*, 'INFOGEST static in vitro simulation of gastrointestinal food digestion', *Nature Protocols 2019 14:4*, vol. 14, no. 4, pp. 991–1014, Mar. 2019, doi: 10.1038/s41596-018-0119-1.
- [50] S. Singh, S. Gamlath, and L. Wakeling, 'Nutritional aspects of food extrusion: a review', *Int J Food Sci Technol*, vol. 42, no. 8, pp. 916–929, Aug. 2007, doi: 10.1111/J.1365-2621.2006.01309.X.
- [51] G. K. Baxevanis *et al.*, 'Tahini consumption improves metabolic and antioxidant status biomarkers in the postprandial state in healthy males', vol. 247, pp. 2721–2728, 2021, doi: 10.1007/s00217-021-03828-5.
- [52] R. Chen, J. Wei, and Y. Gao, 'A review of the study of active components and their pharmacology value in *Lepidium meyenii* (Maca)', *Phytotherapy Research*, vol. 35, no. 12, pp. 6706–6719, Dec. 2021, doi: 10.1002/PTR.7257.
- [53] K. Stamatopoulos, E. Katsoyannos, and A. Chatzilazarou, 'Antioxidant Activity and Thermal Stability of Oleuropein and Related Phenolic Compounds of Olive Leaf Extract after Separation and Concentration by Salting-Out-Assisted Cloud Point Extraction', *Antioxidants 2014, Vol. 3, Pages 229-244*, vol. 3, no. 2, pp. 229–244, Apr. 2014, doi: 10.3390/ANTIOX3020229.
- [54] S. Wang and F. Zhu, 'Chemical composition and health effects of maca (*Lepidium meyenii*)', *Food Chem*, vol. 288, pp. 422–443, Aug. 2019, doi: 10.1016/j.foodchem.2019.02.071.
- [55] L. Wu, M. Zhang, X. Xin, F. Lai, and H. Wu, 'Physicochemical and functional properties of a protein isolate from maca (*Lepidium meyenii*) and the secondary structure and immunomodulatory activity of its major protein component.', *Food Funct*, vol. 10, no. 5, pp. 2894–2905, May 2019, doi: 10.1039/c8fo02490a.
- [56] H. Jemai, A. E. L. Feki, and S. Sayadi, 'Antidiabetic and antioxidant effects of hydroxytyrosol and oleuropein from olive leaves in alloxan-diabetic rats', *J Agric Food Chem*, vol. 57, no. 19, pp. 8798–8804, 2009, doi: 10.1021/JF901280R.
- [57] F. Mariotti, D. Tom , P. Patureau Mirand, F. Ois Mariotti, and D. Tom tom, 'Converting Nitrogen into Protein–Beyond 6.25 and Jones' Factors', *Crit Rev Food Sci Nutr*, vol. 48, no. 2, pp. 177–184, 2008, doi: 10.1080/10408390701279749.
- [58] AOAC Official Methods of Analysis, 'AOAC Official Method 996.01. Fat (total, saturated, unsaturated, and monosaturated) in cereal products: Acid hydrolysis capillary gas chromatographic.', 2000.

- [59] A. Sesmat and J.-F. Meullenet, 'Prediction of Rice Sensory Texture Attributes from a Single Compression Test, Multivariate Regression, and a Stepwise Model Optimization Method', *J Food Sci*, vol. 66, no. 1, 1999, doi: 10.1111/j.1365-2621.2001.tb15593.x.
- [60] J. F. C. Meullenet and J. Gross, 'INSTRUMENTAL SINGLE AND DOUBLE COMPRESSION TESTS TO PREDICT SENSORY TEXTURE CHARACTERISTICS OF FOODS', *J Texture Stud*, vol. 30, no. 2, pp. 167–180, Jul. 1999, doi: 10.1111/J.1745-4603.1999.TB00209.X.
- [61] L. S. Dias-Faceto and A. C. Conti-Silva, 'Texture of extruded breakfast cereals: Effects of adding milk on the texture properties and on the correlations between instrumental and sensory analyses', *J Texture Stud*, vol. 53, no. 2, pp. 220–231, Apr. 2022, doi: 10.1111/JTXS.12666.
- [62] M. Peleg, 'The instrumental texture profile analysis revisited', *J Texture Stud*, vol. 50, no. 5, pp. 362–368, Oct. 2019, doi: 10.1111/JTXS.12392.
- [63] A. Sarkar, F. Kanti, A. Gulotta, B. S. Murray, and S. Zhang, 'Aqueous Lubrication, Structure and Rheological Properties of Whey Protein Microgel Particles', 2017, doi: 10.1021/acs.langmuir.7b03627.
- [64] J. L. Miller and K. L. Watkin, 'The Influence of Bolus Volume and Viscosity on Anterior Lingual Force During the Oral Stage of Swallowing', *Dysphagia*, vol. 11, pp. 117–124, 1996.
- [65] A. Sarkar, K. K. T. Goh, and H. Singh, 'Colloidal stability and interactions of milk-protein-stabilized emulsions in an artificial saliva', *Food Hydrocoll*, vol. 23, no. 5, pp. 1270–1278, Jul. 2009, doi: 10.1016/J.FOODHYD.2008.09.008.
- [66] A. Sarkar, F. Xu, and S. Lee, 'Human saliva and model saliva at bulk to adsorbed phases – similarities and differences', *Adv Colloid Interface Sci*, vol. 273, p. 102034, Nov. 2019, doi: 10.1016/J.CIS.2019.102034.
- [67] H. Stone and J. L. Sidel, 'Affective Testing', *Sensory Evaluation Practices*, pp. 247–277, Jan. 2004, doi: 10.1016/B978-012672690-9/50011-1.
- [68] M. Vandenberghe-Descamps, C. Sulmont-Rossé, C. Septier, G. Feron, and H. Labouré, 'Using food comfortability to compare food's sensory characteristics expectations of elderly people with or without oral health problems', *J Texture Stud*, vol. 48, no. 4, pp. 280–287, Aug. 2017, doi: 10.1111/jtxs.12250.
- [69] H. T. Lawless and H. Heymann, 'SENSORY EVALUATION OF FOOD PRINCIPLES AND PRACTICES'.
- [70] A. Agudelo, P. Varela, and S. Fiszman, 'Methods for a deeper understanding of the sensory perception of fruit fillings', *Food Hydrocoll*, vol. 46, pp. 160–171, Apr. 2015, doi: 10.1016/J.FOODHYD.2014.12.024.
- [71] S. Lee, K. Jo, H. G. Jeong, Y. S. Choi, H. I. Yong, and S. Jung, 'Understanding protein digestion in infants and the elderly: Current in vitro digestion models', *Crit Rev Food Sci Nutr*, 2021, doi: 10.1080/10408398.2021.1957765.
- [72] A. I. Mulet-Cabero *et al.*, 'A standardised semi-dynamic in vitro digestion method suitable for food – an international consensus', *Food Funct*, vol. 11, no. 2, pp. 1702–1720, Feb. 2020, doi: 10.1039/C9FO01293A.
- [73] C. Shani Levi, N. Goldstein, R. Portmann, and U. Lesmes, 'Emulsion and protein degradation in the elderly: Qualitative insights from a study coupling a dynamic in vitro digestion model with proteomic analyses', *Food Hydrocoll*, vol. 69, pp. 393–401, 2017, doi: 10.1016/j.foodhyd.2017.02.017.
- [74] S. David, A. Wojciechowska, R. Portmann, A. Shpigelman, and U. Lesmes, 'The impact of food-grade carrageenans and consumer age on the in vitro proteolysis of whey proteins', *Food Research International*, vol. 130, p. 108964, Apr. 2020, doi: 10.1016/j.foodres.2019.108964.
- [75] M. Ben David, C. Shani Levi, and U. Lesmes, 'Carrageenan impact on digestive proteolysis of meat proteins in meatballs or soluble hydrolyzed collagen', *Food Research International*, vol. 174, p. 113560, Dec. 2023, doi: 10.1016/J.FOODRES.2023.113560.
- [76] C. Shani-Levi *et al.*, 'Extending in vitro digestion models to specific human populations: Perspectives, practical tools and bio-relevant information', *Trends Food Sci Technol*, vol. 60, pp. 52–63, Feb. 2017, doi: 10.1016/J.TIFS.2016.10.017.
- [77] R. Sousa *et al.*, 'In vitro digestibility of dietary proteins and in vitro DIAAS analytical workflow based on the INFOGEST static protocol and its validation with in vivo data', *Food Chem*, vol. 404, p. 134720, Mar. 2023, doi: 10.1016/J.FOODCHEM.2022.134720.
- [78] 'ISO 13903 First edition Animal feeding stuffs–Determination of amino acids content(E) PDF disclaimer(E)', 2005.

- [79] S. Leser, 'The 2013 FAO report on dietary protein quality evaluation in human nutrition: Recommendations and implications', *Nutr Bull*, vol. 38, no. 4, pp. 421–428, Dec. 2013, doi: 10.1111/NBU.12063.
- [80] N. Pineau *et al.*, 'Temporal Dominance of Sensations: Construction of the TDS curves and comparison with time–intensity', *Food Qual Prefer*, vol. 20, no. 6, pp. 450–455, Sep. 2009, doi: 10.1016/J.FOODQUAL.2009.04.005.
- [81] A. S. Levine and J. Ubbink, 'Ultra-processed foods: Processing versus formulation', *Obes Sci Pract*, vol. 9, no. 4, pp. 435–439, Aug. 2023, doi: 10.1002/OSP4.657.
- [82] E. Derbyshire, 'Are all “ultra-processed” foods nutritional demons? A commentary and nutritional profiling analysis.', Dec. 01, 2019, *Elsevier Ltd*. doi: 10.1016/j.tifs.2019.08.023.
- [83] Y. Zhang and E. L. Giovannucci, 'Ultra-processed foods and health: a comprehensive review', *Crit Rev Food Sci Nutr*, vol. 63, no. 31, pp. 10836–10848, 2022, doi: 10.1080/10408398.2022.2084359.
- [84] M. J. Gibney, C. G. Forde, D. Mullally, and E. R. Gibney, 'Ultra-processed foods in human health: A critical appraisal', Sep. 01, 2017, *American Society for Nutrition*. doi: 10.3945/ajcn.117.160440.
- [85] L. A. Tucker and K. S. Thomas, 'Increasing Total Fiber Intake Reduces Risk of Weight and Fat Gains in Women', *J Nutr*, vol. 139, no. 3, pp. 576–581, Mar. 2009, doi: 10.3945/JN.108.096685.
- [86] K. A. Meyer, L. H. Kushi, D. R. Jacobs, J. Slavin, T. A. Sellers, and A. R. Folsom, 'Carbohydrates, dietary fiber, and incident type 2 diabetes in older women', *Am J Clin Nutr*, vol. 71, no. 4, pp. 921–930, Apr. 2000, doi: 10.1093/AJCN/71.4.921.
- [87] Y. Park, L. A. Brinton, A. F. Subar, A. Hollenbeck, and A. Schatzkin, 'Dietary fiber intake and risk of breast cancer in postmenopausal women: the National Institutes of Health–AARP Diet and Health Study', *Am J Clin Nutr*, vol. 90, no. 3, pp. 644–651, Sep. 2009, doi: 10.3945/AJCN.2009.27758.
- [88] M. T. Stroppel, M. C. Ocké, H. C. Boshuizen, F. J. Kok, and D. Kromhout, 'Dietary fiber intake in relation to coronary heart disease and all-cause mortality over 40 y: the Zutphen Study', *Am J Clin Nutr*, vol. 88, no. 4, pp. 1119–1125, Oct. 2008, doi: 10.1093/AJCN/88.4.1119.
- [89] B. Sturtzel, C. Mikulits, C. Gisinger, and I. Elmadfa, 'Use of fiber instead of laxative treatment in a geriatric hospital to improve the wellbeing of seniors', *Journal of Nutrition, Health and Aging*, vol. 13, no. 2, pp. 136–139, Apr. 2009, doi: 10.1007/S12603-009-0020-2/METRICS.
- [90] R. J. Wood *et al.*, 'Effects of a carbohydrate-restricted diet with and without supplemental soluble fiber on plasma low-density lipoprotein cholesterol and other clinical markers of cardiovascular risk', *Metabolism*, vol. 56, no. 1, pp. 58–67, Jan. 2007, doi: 10.1016/J.METABOL.2006.08.021.
- [91] L. Kehoe, J. Walton, and A. Flynn, 'Conference on “Targeted approaches to tackling current nutritional issues” Postgraduate Symposium Nutritional challenges for older adults in Europe: current status and future directions', 2019, doi: 10.1017/S0029665118002744.
- [92] C. Sahu and S. Patel, 'Optimization of maize-millet based soy fortified composite flour for preparation of RTE extruded products using D-optimal mixture design', *J Food Sci Technol*, vol. 58, doi: 10.1007/s13197-020-04771-1.
- [93] R. Photinam and A. Moongngarm, 'Effect of rice flour to corn grits ratio and rice bran oil addition on starch digestibility profiles and properties of extruded multigrain puffed snacks', *Int J Food Sci Technol*, vol. 58, no. 9, pp. 4841–4848, Sep. 2023, doi: 10.1111/IJFS.16598.
- [94] J. P. Singh, A. Kaur, B. Singh, N. Singh, and B. Singh, 'Physicochemical evaluation of corn extrudates containing varying buckwheat flour levels prepared at various extrusion temperatures', *J Food Sci Technol*, vol. 56, no. 4, pp. 2205–2212, Apr. 2019, doi: 10.1007/S13197-019-03703-Y/TABLES/2.
- [95] R. Kadowaki, H. Kimura, and N. Inou, 'New Estimation Methods of Young's Modulus and Rupture Strength of Snack Foods Based on Microstructure', *J Texture Stud*, vol. 47, no. 1, pp. 3–13, Feb. 2016, doi: 10.1111/JTXS.12154.
- [96] M. Michalak-Majewska *et al.*, 'COMPARATIVE ANALYSIS OF SELECTED PHYSICO-CHEMICAL AND TEXTURAL PROPERTIES OF BREAD SUBSTITUTES', *Acta Universitatis Cibiniensis Series E: FOOD TECHNOLOGY*, vol. 99, no. 1, doi: 10.2478/auaft-2020-0009.
- [97] H. H. Kara, A. Araiza-Calahorra, N. M. Rigby, and A. Sarkar, 'Flaxseed oleosomes: Responsiveness to physicochemical stresses, tribological shear and storage', *Food Chem*, vol. 431, p. 137160, Jan. 2024, doi: 10.1016/J.FOODCHEM.2023.137160.

- [98] M. E. Malone, I. A. M. Appelqvist, and I. T. Norton, 'Oral behaviour of food hydrocolloids and emulsions. Part 1. Lubrication and deposition considerations', doi: 10.1016/S0268-005X(03)00097-3.
- [99] J. Chen, N. Khandelwal, Z. Liu, and T. Funami, 'Influences of food hardness on the particle size distribution of food boluses', *Arch Oral Biol*, vol. 58, no. 3, pp. 293–298, Mar. 2013, doi: 10.1016/J.ARCHORALBIO.2012.10.014.
- [100] M. Vandenberghe-Descamps, H. Labouré, C. Septier, G. Feron, and C. Sulmont-Rossé, 'Oral comfort: A new concept to understand elderly people's expectations in terms of food sensory characteristics', *Food Qual Prefer*, vol. 70, pp. 57–67, Dec. 2018, doi: 10.1016/J.FOODQUAL.2017.08.009.
- [101] D. S. Khamrova and S. V. Popov, 'A secret of salivary secretions: Multimodal effect of saliva in sensory perception of food', *Eur J Oral Sci*, vol. 130, no. 2, p. e12846, Apr. 2022, doi: 10.1111/EOS.12846.
- [102] H. Chen, M. Iinuma, M. Onozuka, and K. Y. Kubo, 'Chewing Maintains Hippocampus-Dependent Cognitive Function', *Int J Med Sci*, vol. 12, no. 6, p. 502, 2015, doi: 10.7150/IJMS.11911.
- [103] F. Barbé *et al.*, 'Acid and rennet gels exhibit strong differences in the kinetics of milk protein digestion and amino acid bioavailability', *Food Chem*, vol. 143, pp. 1–8, Jan. 2014, doi: 10.1016/J.FOODCHEM.2013.07.100.
- [104] K. Nyemb *et al.*, 'The structural properties of egg white gels impact the extent of in vitro protein digestion and the nature of peptides generated', *Food Hydrocoll*, vol. 54, pp. 315–327, Mar. 2016, doi: 10.1016/J.FOODHYD.2015.10.011.
- [105] D. Dupont *et al.*, 'Can dynamic *in vitro* digestion systems mimic the physiological reality?', *Crit Rev Food Sci Nutr*, vol. 59, no. 10, pp. 1546–1562, May 2019, doi: 10.1080/10408398.2017.1421900.
- [106] D. Duijsens *et al.*, 'Strategic choices for in vitro food digestion methodologies enabling food digestion design', vol. 126, pp. 61–72, Aug. 2022, doi: 10.1016/J.TIFS.2022.06.017.
- [107] A. Mackie and A. Macierzanka, 'Colloidal aspects of protein digestion', *Curr Opin Colloid Interface Sci*, vol. 15, no. 1–2, pp. 102–108, 2010, doi: 10.1016/j.cocis.2009.11.005.
- [108] R. Sousa, R. Portmann, S. Dubois, I. Recio, and L. Egger, 'Protein digestion of different protein sources using the INFOGEST static digestion model', *Food Research International*, vol. 130, p. 108996, Apr. 2020, doi: 10.1016/J.FOODRES.2020.108996.
- [109] S. A. Adibi and D. W. Mercer, 'Protein Digestion in Human Intestine as Reflected in Luminal, Mucosal, and Plasma Amino-Acid Concentrations after Meals', *Journal of Clinical Investigation*, vol. 52, no. 7, pp. 1586–1594, 1973, doi: 10.1172/Jci107335.
- [110] Roger. Lentle and P. W. M. Janssen, 'The physical processes of digestion', 2011, Accessed: Jan. 26, 2022. [Online]. Available: https://books.google.com/books/about/The_Physical_Processes_of_Digestion.html?hl=iw&id=kN1edFo-1bUC
- [111] S. M. Loveday, 'Food Proteins: Technological, Nutritional, and Sustainability Attributes of Traditional and Emerging Proteins', Mar. 25, 2019, *Annual Reviews Inc.* doi: 10.1146/annurev-food-032818-121128.
- [112] J. K. Mathai, Y. Liu, and H. H. Stein, 'Values for digestible indispensable amino acid scores (DIAAS) for some dairy and plant proteins may better describe protein quality than values calculated using the concept for protein digestibility-corrected amino acid scores (PDCAAS)', *British Journal of Nutrition*, vol. 117, no. 04, pp. 490–499, Feb. 2017, doi: 10.1017/S0007114517000125.
- [113] M. S. Alam, J. Kaur, H. Khaira, and K. Gupta, 'Extrusion and Extruded Products: Changes in Quality Attributes as Affected by Extrusion Process Parameters: A Review', *Crit Rev Food Sci Nutr*, vol. 56, no. 3, pp. 445–473, Feb. 2016, doi: 10.1080/10408398.2013.779568.