

## **The potential of transgenic legumes in integrated bruchid management: assessing the impact on bruchid parasitoids**

Lüthi, C.\*#, Álvarez-Alfageme, F., Romeis, J.

Agroscope Reckenholz-Tänikon Research Station ART, Reckenholzstrasse 191, 8046 Zurich, Switzerland.

Email: christoph.luethi@art.admin.ch

\* Corresponding author

# Presenting author

DOI: 10.5073/jka.2010.425.167.251

### **Abstract**

Leguminous seeds are an important staple food and source of nutrition in many countries. Bruchid beetles (Coleoptera: Bruchidae) are responsible for the greatest post-harvest losses to stored legumes. A powerful strategy to control bruchid infestations is the combination of plant resistance factors and biological control provided by parasitoids. Potent resistance factors are  $\alpha$ -amylase inhibitors ( $\alpha$ AI) which inhibit the starch metabolism in sensitive insects. Genetic engineering has been used to transfer  $\alpha$ AI-1 from the common bean (*Phaseolus vulgaris*) to other leguminous plants which are subsequently protected from the attack by several bruchid species. However, there are concerns regarding the effects that the expressed insecticidal protein might have on non-target organisms. Here, we present an approach to assess the impact of  $\alpha$ AI-1 genetically modified legumes on bruchid parasitoids.

**Keywords:** Risk assessment, Genetically modified plants, Non-target organisms;  $\alpha$ -amylase inhibitor;  $\alpha$ AI-1

### **1. Introduction**

Legume seeds are an important source of nutrition for both humans and livestock. Their seeds are rich in proteins, carbohydrates and lipids, and they can be stored over extended periods. Additionally, the nitrogen-fixing abilities of the plants are important for the management of soil fertility. All these properties match perfectly with the requirements of small-scale, low-income farmers in developing countries.

Bruchid beetles (Coleoptera: Bruchidae) are responsible for the largest post-harvest losses to stored seeds, directly through consumption of the resource and, secondarily, through the qualitative deterioration of the commodity or the reduced stock viability. The females lay their eggs on the seed surface and the larvae burrow into the seed, where they feed and complete their development (Southgate, 1979). The beetles usually continue to multiply during seed storage, which can lead to extensive or even total losses, especially if the seeds are stored for long periods. Surface and fumigant chemical applications are thought to be the most effective methods for managing bruchid infestations. However, prohibitive costs, which limit their application to large scale or extended storage, and the risks of adverse secondary effects from such treatments, have driven the exploration of alternative strategies to manage bruchid infestations. These include biological control (Sanon et al., 1998; Gauthier et al., 1999; Schmale et al., 2006) and plant resistance factors (Ignacimuthu et al., 2000; Schmale et al., 2003; Appleby and Credland, 2004).

A crop protection tool with high potential for small-scale farmers is genetic engineering. Although the largest areas of genetically modified (GM) crop production have been in industrial countries, it's the small-scale farmers in developing countries that might benefit the most from this technology (Wambugu, 1999; Thomson, 2008). In 2008, 90% of the farmers planting GM crops were in developing countries (James, 2008). To date, all commercially cultivated insect-resistant GM crops are expressing Cry proteins derived from the soil bacterium *Bacillus thuringiensis* (Berliner) (so called *Bt* crops). The potential of insects to evolve resistance against the deployed *Bt* Cry toxins, their narrow spectrum of activity and the risk of infringing existing patents have driven the development of alternative insecticidal traits for genetic engineering, including inhibitors of digestive enzymes (Malone et al., 2008).

## 2. Using a *Phaseolus vulgaris* resistance factor for genetic engineering

The common bean, *Phaseolus vulgaris* L. (Fabaceae) and other *Phaseolus* species possess a family of evolutionary related defence proteins including phytohemagglutinin (PHA), arcelin (Arc) and  $\alpha$ -amylase inhibitors ( $\alpha$ AI) (Chrispeels and Raikhel, 1991). The genes of these three proteins are encoded by a single locus in the *P. vulgaris* genome (Nodari et al., 1993) and it is likely that these homologous genes have arisen by duplication of an ancestral gene. The proteins feature different modes of action and insecticidal properties against bruchids (Leavitt et al., 1977; Liener, 1986; Osborn et al., 1988; Janarthan et al., 2008; Velten et al., 2007b).  $\alpha$ AI can strongly inhibit the activity of  $\alpha$ -amylases. These enzymes hydrolyse starch or glycogen and play a key role in the carbohydrate metabolism of microorganisms, plants and animals. Several insects, especially those feeding on starchy seeds during any period of their life cycle like bruchid beetles, depend on  $\alpha$ -amylases for survival (Grossi de Sa et al., 1997; Franco et al., 2002). This feature has attracted notice to  $\alpha$ AI, making them a promising candidate for genetic engineering. The  $\alpha$ AI of *P. vulgaris* exists in at least two different allelic variants. The isoform found in cultivated beans is called  $\alpha$ AI-1 (Moreno and Chrispeels, 1989); a second variant,  $\alpha$ AI-2, is found in some wild accessions of the common bean that contain Arc as the major storage protein instead of phaseolin (Suzuki et al., 1993). Of particular interest is that the two isoforms differ in their specificity towards  $\alpha$ -amylases. With the exception of one major storage pest, *Acanthoscelides obtectus* (Say) (Coleoptera: Bruchidae), one or both  $\alpha$ AI from the common bean have been shown to inhibit midgut amylases of major bruchid pests found worldwide (Ishimoto and Kitamura, 1992; Franco et al., 2002) (Table 1). The successful transfer of common bean  $\alpha$ AI into tobacco plants (Altabella and Chrispeels, 1990) indicated that they could also be engineered to other plant species. Subsequently, genes encoding  $\alpha$ AI-1 and/or  $\alpha$ AI-2 from *P. vulgaris* were introduced by methods of genetic engineering into peas (*Pisum sativum* L.), cowpeas (*Vigna unguiculata* L.), azuki beans (*Vigna angularis* (Willdenow)) and chickpeas (*Cicer arietinum* L.) (Shade et al., 1994; Schroeder et al., 1995; Ishimoto et al., 1996; Morton et al., 2000; Solleti et al., 2008). The  $\alpha$ AI gene construct introduced in these plants is regulated by flanking sequences from the seed-specific bean phytohemagglutinin PHA (*dlec*) gene. This promoter regulates the expression of the  $\alpha$ AI restrictively to the cotyledon and embryonic axis of the developing seed (Schroeder et al., 1995), targeting exclusively seed-feeding herbivores like bruchids.

The potential of GM legumes expressing  $\alpha$ AI to control bruchids has been confirmed in several studies. In peas, levels of expression of  $\alpha$ AI ranged between 1.5 and 3.5% of total soluble protein, providing 100% control of *Bruchus pisorum* (L.) under glasshouse (Schroeder et al., 1995) and field conditions (Morton et al., 2000). Azuki beans expressing 0.9%  $\alpha$ AI-1 (per dry weight) provided 100% control of the two bruchids, *Callosobruchus chinensis* L. and *Callosobruchus maculatus* F., both important pest species in East Asia (Ishimoto et al., 1996). Similarly,  $\alpha$ AI-1-GM cowpeas and chickpeas have been reported to strongly inhibit the development of *C. chinensis* and *C. maculatus* (Sarmah et al., 2004; Solleti et al., 2008).


## 3. Compatibility of $\alpha$ AI with biocontrol agents

The potential of combining plant resistance factors together with biological control agents, especially parasitoids, has been shown to be a powerful method to control storage pests like bruchids (Schmale et al., 2003; Velten et al., 2008). One major concern regarding the use of insect-resistant GM plants is the effect that the expressed insecticidal protein might have on non-target organisms. This is especially relevant for traits with a broad spectrum of activity such as  $\alpha$ AI where any organism relying on  $\alpha$ -amylases for carbohydrate digestion is potentially affected by the inhibitor. The potential impact of  $\alpha$ AI on bruchid parasitoids has never been evaluated. Kluh et al. (2005) extensively screened the inhibitory activity of  $\alpha$ AI-1 against 24 insect species from eight different orders. The most sensitive species belonged to the orders of Coleoptera, Diptera and Hymenoptera. The Hymenoptera tested included an endoparasitic wasp, *Venturia canescens* (Gravenhorst) (Hymenoptera: Ichneumonidae). However, no bruchid parasitoids were included and whether they rely on  $\alpha$ -amylases is still unknown.

## 4. Assessing the impact of legumes expressing $\alpha$ AI-1 on bruchid parasitoids

Environmental risk assessment for non-target organisms is a required step in the evaluation process of new GM plants. This is particularly true for GM crops expressing insecticidal proteins. The assessment typically follows a tiered framework which is conceptually similar to that of pesticides (Hill and Sendashonga, 2003; Romeis et al., 2008).

In an initial step one has to define what we do not want to see harmed by the GM trait. In the case of  $\alpha$ AI-1 expressing legume seeds this is the biological control function provided by parasitic wasps that contribute to the control of bruchid species which are not or insufficiently affected by the introduced insecticidal trait. We can then develop a conceptual model describing a pathway how the presence of  $\alpha$ AI-1 in the GM seeds can cause harm to biological control (Fig. 1). Following the different steps of the conceptual model allows a comprehensive evaluation of the risk for the parasitoids.



**Figure 1** Pathway on how  $\alpha$ AI-1 from *P. vulgaris* expressed in transgenic legume seeds could harm the biological control function provided by bruchid parasitoids.

The most basic requirement to make bruchid parasitoids a potential target of  $\alpha$ AI-1 is the presence of  $\alpha$ -amylases. Therefore the  $\alpha$ -amylase activity was first characterized in five different parasitic wasps which are commonly used to control bruchids (Step 1): *Anisopteromalus calandrae* (Howard) (Pteromalidae), *Dinarmus basalis* (Rondani) (Pteromalidae), *Lariophagus distinguendus* (Förster) (Pteromalidae), *Eupelmus vuillei* (Crawford) (Eupelmidae) and *Heterospilus prosopidis* (Viereck) (Braconidae). The characterization of such enzyme activity was based on the *in vitro* characteristics of complete insect extracts. Larval and female extracts of all species were able to hydrolyze the specific substrate potato starch although a higher activity was observed in the latter. Moreover, all extracts were highly susceptible to the specific inhibitors acarbose and wheat  $\alpha$ AI. Taken all together, these results suggest that both larvae and females rely on  $\alpha$ -amylase activity for carbohydrate digestion. Once  $\alpha$ -amylase activity was detected, we went to the second step. The *in vitro* susceptibility to  $\alpha$ AI-1 was determined by using different concentrations of the inhibitor in order to construct dose-response curves for all species. The  $\alpha$ -amylase activities from all larval and female extracts were highly susceptible to  $\alpha$ AI-1. This finding necessitates testing the impact of  $\alpha$ AI-1 on the parasitoids *in vivo*.

The subsequent steps of the assessment will include tritrophic experiments using different plant-host-parasitoid systems. We selected  $\alpha$ AI-1 expressing GM cowpea and chickpea lines and three bruchid species as hosts for the parasitoids, namely *C. maculatus*, *C. chinensis* and *A. obtectus*. The two former are reported being susceptible, the latter being resistant to  $\alpha$ AI-1 (Table 1). Bioassays with bruchids developing in GM cowpea and chickpea lines, their non-transformed isolines and other non-GM varieties will be performed to assess the variance of resistance. Measuring different life-history traits will allow verifying the susceptibility of the bruchids to  $\alpha$ AI-1 reported in the literature. The exposure to the inhibitor at the third trophic level will also be investigated; using Western Blotting or ELISA tests, the presence of the insecticidal compound in the body of collected bruchid larvae as well as parasitoid larvae and adult females of host-feeding parasitoids will be analyzed. These data allow retracing the path of  $\alpha$ AI-1 through the food chain and help to understand the cause of possible impacts on the non-target organisms (Step 3).

**Table 1** Present known distribution of major bruchid pests and their sensitivity to  $\alpha$ AI-1 and  $\alpha$ AI-2 from *P. vulgaris*.

Species	Distribution Worldwide						Inhibition of midgut $\alpha$ -amylase	
	North America	South America	Africa	Asia	Australia	Europe	$\alpha$ AI-1	$\alpha$ AI-2
<i>Acanthoscelides obtectus</i>	(+)	(+)	+	+	+	+	n <sup>3</sup>	n <sup>3</sup>
<i>Bruchus pisorum</i>	+	+	+	(+)	+	+	y <sup>2</sup>	p <sup>2</sup>
<i>Callosobruchus analis</i>	+		+	(+)			y <sup>2</sup>	
<i>Callosobruchus chinensis</i>	+	+	(+)	(+)	+	+	y <sup>1,2,3</sup>	n <sup>1,3</sup>
<i>Callosobruchus maculatus</i>	+	+	(+)	(+)	+	+	y <sup>1,2</sup>	y <sup>3</sup>
<i>Zabrotes subfasciatus</i>	(+)	(+)	+	+		+	n <sup>1,2,3</sup>	y <sup>1,2,3</sup>

(+) indigenous; + established; y = yes; n = no; p = partial; <sup>1</sup> artificial diet; <sup>2</sup> transgenic seeds; <sup>3</sup> amylase inhibitory activity

In the fourth step, tritrophic studies will be conducted to establish the host-mediated effect of  $\alpha$ AI-1 on the selected susceptible parasitoids. Parasitoid females will be provided GM seeds expressing  $\alpha$ AI-1 or untransformed control seeds infested with larvae of susceptible or tolerant bruchid species and parasitoid performance (e.g., survival, fecundity) will be assessed according to Velten et al. (2007a). If detrimental effects at the third trophic level are observed, bruchid-parasitoid population dynamic experiments will be conducted in the final step to assess long-term effects on the non-target population (Wäckers, 2003).

## 5. Conclusions

GM legumes expressing  $\alpha$ AI-1 from *P. vulgaris* are protected from the attack by major bruchid pests. Some species, including *A. obtectus* and *Zabrotes subfasciatus* (Boheman), however, remain unaffected by this resistant factor. It is thus important that the biological control of these species is not disrupted by the insecticidal trait expressed in the GM legumes. However, the broad range of activity of this inhibitor and its possible transfer through the food-chain necessitate a detailed analysis of the possible impacts on beneficial non-target organisms. *In vitro* inhibition studies showed that  $\alpha$ AI-1 inhibits  $\alpha$ -amylase activity in larvae and females of several species of bruchid parasitoids. Consequently, parasitoids might be adversely affected when developing in or feeding on bruchid pests that are not controlled by the GM trait. The assessment of these non-target effects is necessary to ensure the compatible use of biological control and the GM host plant resistance trait for a sustainable control of bruchids.

## References

- Altabella, T., Chrispeels, M.J., 1990. Tobacco plants transformed with the bean  $\alpha$ AI gene express an inhibitor of insect  $\alpha$ -amylase in their seeds. *Plant Physiology* 93, 805-810.
- Appleby, J.H., Credland, P.F., 2004. Environmental conditions affect the response of West African *Callosobruchus maculatus* (Coleoptera: Bruchidae) populations to susceptible and resistant cowpeas. *Journal of Stored Products Research* 40, 269-287.
- Chrispeels, M.J., Raikhel, N.V., 1991. Lectins, lectin genes, and their role in plant defense. *Plant Cell* 3, 1-9.
- Franco, O.L., Rigden, D.J., Melo, F.R., Grossi de Sa, M.F., 2002. Plant alpha-amylase inhibitors and their interaction with insect alpha-amylases - structure, function and potential for crop protection. *European Journal of Biochemistry* 269, 397-412.
- Gauthier, N., Sanon, A., Monge, J.P., Huignard, J., 1999. Interspecific relations between two sympatric species of hymenoptera, *Dinarmus basalis* (Rond) and *Eupelmus vuilleti* (Crw), ectoparasitoids of the bruchid *Callosobruchus maculatus* (F.). *Journal of Insect Behavior* 12, 399-413.
- Grossi de Sa, M.F., Mirkov, T.E., Ishimoto, M., Colucci, G., Bateman, K.S., Chrispeels, M.J., 1997. Molecular characterization of a bean  $\alpha$ -amylase inhibitor that inhibits the  $\alpha$ -amylase of the Mexican bean weevil *Zabrotes subfasciatus*. *Planta* 203, 295-303.
- Hill, R.A., Sendashonga, C., 2003. General principles for risk assessment of living modified organisms: Lessons from chemical risk assessment. *Environmental Biosafety Research* 2, 81-88.
- Ignacimuthu, S., Janarthanan, S., Balachandran, B., 2000. Chemical basis of resistance in pulses to *Callosobruchus maculatus* (F.) (Coleoptera: Bruchidae). *Journal of Stored Products Research* 36, 89-99.
- Ishimoto, M., Kitamura, K., 1992. Tolerance to the seed  $\alpha$ -amylase inhibitor by the two insect pests of the common bean, *Zabrotes subfasciatus* and *Acanthoscelides obtectus* (Coleoptera: Bruchidae). *Applied Entomology and Zoology* 27, 243-251.
- Ishimoto, M., Sato, T., Chrispeels, M.J., Kitamura, K., 1996. Bruchid resistance of transgenic azuki bean expressing seed  $\alpha$ -amylase inhibitor of common bean. *Entomologia Experimentalis et Applicata* 79, 309-315.
- James, C., 2008. Global Status of Commercialized Biotech/GM Crops: 2008. ISAAA Brief No. 39, 1-20.
- Janarthanan, S., Suresh, P., Radke, G., Morgan, T.D., Oppert, B., 2008. Arcelins from an Indian wild pulse, *Lablab purpureus*, and insecticidal activity in storage pests. *Journal of Agricultural and Food Chemistry* 56, 1676-1682.
- Kluh, I., Horn, M., Hýblová, J., Hubert, J., Dolecková-Maresová, L., Voburka, Z., Kudlíková, I., Kocourek, F., Mares, M., 2005. Inhibitory specificity and insecticidal selectivity of  $\alpha$ -amylase inhibitor from *Phaseolus vulgaris*. *Phytochemistry* 66, 31-39.
- Leavitt, R.D., Felsted, R.L., Bachur, N.R., 1977. Biological and biochemical properties of *Phaseolus vulgaris* isolecithins. *Journal of Biological Chemistry* 252, 2961-2966.
- Liener, I. E., 1986. Nutritional significance of lectins in the diet. In: Liener, I.E., Sharon, N., Goldstein, I.J. (Eds), *The Lectins*. Academic Press, San Diego, CA, USA, pp. 527-552.
- Malone, L.A., Gatehouse, A.M.R., Barratt, B.I.P., 2008. Beyond *Bt*: alternative strategies for insect-resistant genetically modified crops. In: Romeis, J., Shelton, A.M., Kennedy, G.G. (Eds), *Integration of insect-resistant genetically modified crops within IPM programs*. Springer Science + Business Media B.V., Dordrecht, The Netherlands, pp. 357-417.
- Moreno, J., Chrispeels, M.J., 1989. A lectin gene encodes the  $\alpha$ -amylase inhibitor of the common bean. *Proceedings of the National Academy of Sciences of the United States of America* 86, 7885-7889.
- Morton, R.L., Schroeder, H.E., Bateman, K.S., Chrispeels, M.J., Armstrong, E., Higgins, T.J.V., 2000. Bean  $\alpha$ -amylase inhibitor 1 in transgenic peas (*Pisum sativum*) provides complete protection from pea weevil (*Bruchus pisorum*) under field conditions. *Proceedings of the National Academy of Sciences of the United States of America* 97, 3820-3825.
- Nodari, R.O., Tsail, S.M., Gilbertson, R.L., Gepts, P., 1993. Towards an integrated linkage map of common bean 2. Development of an RFLP-based linkage map. *Theoretical and Applied Genetics* 85, 513-520.
- Osborn, T.C., Alexander, D.C., Sun, S.S.M., Cardona, C., Bliss, F.A., 1988. Insecticidal activity and lectin homology of arcelin seed protein. *Science* 240, 207-210.
- Romeis, J., Bartsch, D., Bigler, F., Candolfi, M.P., Gielkens, M.M.C., Hartley, S.E., Hellmich, R.L., Huesing, J.E., Jepson, P.C., Layton, R., Quemada, H., Raybould, A., Rose, R.I., Schiemann, J., Sears, M.K., Shelton, A.M., Sweet, J., Vaituzis, Z., Wolt, J.D., 2008. Assessment of risk of insect-resistant transgenic crops to nontarget arthropods. *Nature Biotechnology* 26, 203-208.

- Sanon, A., Ouedraogo, A.P., Tricault, Y., Credland, P.F., Huignard, J., 1998. Biological control of bruchids in cowpea stores by release of *Dinarmus basalis* (Hymenoptera: Pteromalidae) adults. *Environmental Entomology* 27, 717-725.
- Sarmah, B.K., Moore, A., Tate, W., Molvig, L., Morton, R.L., Rees, D.P., Chiaiese, P., Chrispeels, M.J., Tabe, L.M., Higgins, T.J.V., 2004. Transgenic chickpea seeds expressing high levels of a bean  $\alpha$ -amylase inhibitor. *Molecular Breeding* 14, 73-82.
- Schmale, I., Wäckers, F.L., Cardona, C., Dorn, S., 2003. Combining parasitoids and plant resistance for the control of the bruchid *Acanthoscelides obtectus* in stored beans. *Journal of Stored Products Research* 39, 401-411.
- Schmale, I., Wäckers, F.L., Cardona, C., Dorn, S., 2006. Biological control of the bean weevil, *Acanthoscelides obtectus* (Say) (Col.: Bruchidae), by the native parasitoid *Dinarmus basalis* (Rondani) (Hym.: Pteromalidae) on small-scale farms in Colombia. *Journal of Stored Products Research* 42, 31-41.
- Schroeder, H.E., Gollasch, S., Moore, A., Tabe, L.M., Craig, S., Hardie, D.C., Chrispeels, M.J., Spencer, D., Higgins, T.J.V., 1995. Bean  $\alpha$ -amylase inhibitor confers resistance to the pea weevil (*Bruchus pisorum*) in transgenic peas (*Pisum sativum* L.). *Plant Physiology* 107, 1233-1239.
- Shade, R.E., Schroeder, H.E., Pueyo, J.J., Tabe, L.M., Murdock, L.L., Higgins, T.J.V., Chrispeels, M.J., 1994. Transgenic pea seeds expressing the  $\alpha$ -amylase inhibitor of the common bean are resistant to bruchid beetles. *Nature Biotechnology* 12, 793-796.
- Solleti, S.K., Bakshi, S., Purkayastha, J., Panda, S.K., Sahoo, L., 2008. Transgenic cowpea (*Vigna unguiculata*) seeds expressing a bean  $\alpha$ -amylase inhibitor 1 confer resistance to storage pests, bruchid beetles. *Plant Cell Reports* 27, 1841-1850.
- Southgate, B.J., 1979. Biology of the Bruchidae. *Annual Review of Entomology* 24, 449-473.
- Suzuki, K., Ishimoto, M., Kikuchi, F., Kitamura, K., 1993. Growth-inhibitory effect of an  $\alpha$ -amylase inhibitor from the wild common bean resistant to the Mexican bean weevil (*Zabrotes subfasciatus*). *Japanese Journal of Breeding* 43, 257-265.
- Thomson, J.A., 2008. The role of biotechnology for agricultural sustainability in Africa. *Philosophical Transactions of the Royal Society B - Biological Sciences* 363, 905-913.
- Velten, G., Rott, A.S., Cardona, C., Dorn, S., 2007a. Effects of a plant resistance protein on parasitism of the common bean bruchid *Acanthoscelides obtectus* (Coleoptera: Bruchidae) by its natural enemy *Dinarmus basalis* (Hymenoptera: Pteromalidae). *Biological Control* 43, 78-84.
- Velten, G., Rott, A.S., Cardona, C., Dorn, S., 2007b. The inhibitory effect of the natural seed storage protein arcelin on the development of *Acanthoscelides obtectus*. *Journal of Stored Products Research* 43, 550-557.
- Velten, G., Rott, A.S., Conde Petit, B.J., Cardona, C., Dorn, S., 2008. Improved bruchid management through favorable host plant traits and natural enemies. *Biological Control* 47, 133-140.
- Wäckers, F.L., 2003. The effect of food supplements on parasitoid-host dynamics. In: Van Driesche, R.G. (Ed), *Proceedings of the First International Symposium on Biological Control of Arthropods*, 14-18 January, 2002, Honolulu, HI, USA. US Department of Agriculture, Forest Service, Morgantown, WV, USA, pp. 226-231.
- Wambugu, F., 1999. Why Africa needs agricultural biotech. *Nature* 400, 15-16.