



# The weakest link: Haploid honey bees are more susceptible to neonicotinoid insecticides

Andrea Friedli<sup>a</sup>, Geoffrey R. Williams<sup>b</sup>, Selina Bruckner<sup>b</sup>, Peter Neumann<sup>a, c</sup>, Lars Straub<sup>a, c, \*</sup>

<sup>a</sup> Institute of Bee Health, Vetsuisse Faculty, University of Bern, Bern, Switzerland

<sup>b</sup> Department of Entomology and Plant Pathology, Auburn University, Auburn, AL, USA

<sup>c</sup> Agroscope, Swiss Bee Research Centre, Bern, Switzerland

## HIGHLIGHTS

- Neonicotinoid insecticides affected the developmental stability of honey bees.
- Haploid males were more susceptible to neonicotinoids than diploid females.
- Heterozygosity may play a key role in buffering insecticide exposure.

## ARTICLE INFO

### Article history:

Received 6 August 2019

Received in revised form

17 October 2019

Accepted 17 October 2019

Available online 22 October 2019

Handling Editor: Willie Peijnenburg

### Keywords:

*Apis mellifera*

Xenobiotics

Fluctuating asymmetry

Wing venation

Haplodiploidy

Sublethal

## ABSTRACT

Neonicotinoid insecticides are currently of major concern for the health of wild and managed insects that provide key ecosystem services like pollination. Even though sublethal effects of neonicotinoids are well known, there is surprisingly little information on how they possibly impact developmental stability, and to what extent genetics are involved. This holds especially true for haploid individuals because they are hemizygous at detoxification loci and may be more susceptible. Here we take advantage of haplodiploidy in Western honey bees, *Apis mellifera*, to show for the first time that neonicotinoids affect developmental stability in diploid females (workers), and that haploid males (drones) are even more susceptible. Phenotypic fore wing venation abnormalities and fluctuating wing asymmetry, as measures of developmental instability, were significantly increased under field-realistic neonicotinoid-exposure of colonies. The higher susceptibility of haploid drones suggests that heterozygosity can play a key role in the ability to buffer the sublethal effects of neonicotinoids. Aiming to improve conservation efforts, our findings highlight the urgent need to better understand the role that genetics plays at enabling non-target organisms to cope with insecticide exposure.

© 2019 The Authors. Published by Elsevier Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

## 1. Introduction

Insects are essential for the functionality of terrestrial ecosystems (Hallmann et al. 2017), and also play an important role in human food security such as by pollination of crops (Garibaldi et al. 2016). Therefore, the increasing global application of insecticides (Schreinemachers and Tipraqsa, 2012) together with major declines of wild and losses of managed insect species (Neumann and Carreck, 2010; Hallmann et al. 2017) has raised considerable

concern over the stability of the ecosystem services provided by non-target insects (Potts et al. 2016). Neonicotinoids currently represent the most commonly employed insecticides in agricultural pest management practices (Simon-Delso et al. 2015). These insecticides act as antagonists of the nicotinic acetylcholine receptors, which leads to paralysis and ultimately death of the organism (Blacquière et al. 2012). Several studies have shown that they can induce sublethal effects on various invertebrates (Neumann et al. 2015), such as reducing fitness in solitary bees (Sandrock et al. 2014a), impairing bumble bee crop pollination services (Stanley et al. 2015), and comprising reproductive capacity of honey bee, queens and drones (Williams et al. 2015; Straub et al. 2016). Nevertheless, there is still an ongoing debate on the role of

\* Corresponding author. Institute of Bee Health Vetsuisse Faculty, University of Bern Schwarzenburgstrasse, 1613097, Bern, Switzerland.

E-mail address: [lars.straub@vetsuisse.unibe.ch](mailto:lars.straub@vetsuisse.unibe.ch) (L. Straub).

neonicotinoids for the decline of wild insect populations and losses of managed ones (Goulson, 2013; Carreck and Ratnieks, 2014), creating demand for an even better understanding of the sublethal effects of these agrochemicals. Even though the negative effects of other insecticides on morphogenetic development have been shown (Prado-Silva et al. 2018), there is surprisingly very little understanding of the potential effects of neonicotinoid insecticides on ontogenetic developmental stability.

Developmental stability is the ability of an organism to produce a consistent phenotype under any given environmental condition (Klingenberg, 2019). Disruptions in development can have broad consequences (Willmore and Hallgrímsson, 2005), including reduced fitness (Møller, 1997). Fluctuating asymmetry, described as random deviations from perfect asymmetry in bilateral traits, as well as the frequency of phenotypic abnormalities (e.g. wing venation abnormalities), are two common parameters used to measure developmental precision in insects (Lopuch and Tofilski, 2016). Consequently, increased fluctuating asymmetry and phenotypic abnormalities suggest impaired developmental stability (Klingenberg, 2019), and are regarded as surrogates for the health of insect populations (Beasley et al. 2013). Such developmental instability may be attributed to the disability of an individual to buffer against both genetic (e.g. mutations) or environmental factors (e.g. parasites, toxins) during morphogenesis (Müller et al. 2017; Gerard et al. 2018).

Past studies have shown that various stressors can cause fluctuating wing asymmetry, such as insecticides or pathogens (Chang et al. 2007; Gerard et al. 2018). Asymmetries in morphological structures imply perturbation in developmental homeostasis at the molecular, chromosomal and epigenetic levels, but underlying mechanisms are unknown (Klingenberg and Nijhout, 1999). It is believed that genetic factors and physiological changes may play a crucial role in the ability to reach developmental stability (Chang et al. 2007), which includes managing xenobiotics by means of detoxifying enzymes (Derecka et al. 2013; Abbo et al. 2016). This may be particularly apparent in haploid individuals that are hemizygous at detoxification loci (O'Donnell and Beshers, 2004) because heterozygosity likely enhances resistance towards parasitic and pathogenic stress (Baer and Schmid-Hempel, 2003). Indeed, haploid hymenopterans were more susceptible when compared to their diploid counterparts under pathogen challenge (Retschnig et al. 2014; Strobl et al. 2019). Moreover, in various insect orders (Hymenoptera, Thysanoptera, and Coleoptera), males usually develop from unfertilized eggs and are therefore haploid (Evans et al. 2004), whereas females are diploid (Ross et al. 2019). In light of recent major declines of insects (Hallmann et al. 2017), the possible higher susceptibility of haploid male insects may constitute a currently neglected gap in our knowledge despite the ample body of literature on insecticides. In particular, the possible role of heterozygosity for the developmental stability under neonicotinoid exposure is currently unknown.

For the first time, we compared susceptibility between the haploid male drone and diploid female worker honey bees to neonicotinoid insecticides by evaluating effects on developmental stability. The western honey bee (*Apis mellifera*) has historically served as a model organism to investigate the effects of environmental and anthropogenic stress on biological organisms, mainly because of their role as a keystone pollinator of agricultural and wild plants (Calderone, 2012), as well as their well-known biology (EFSA, 2014). We compared the frequency of phenotypic fore wing venation abnormalities and the degree of fluctuating fore wing asymmetry between diploid female worker and haploid male honey bees. Colonies were exposed to chronic field-realistic concentrations of two commonly applied neonicotinoid insecticides –

thiamethoxam and its primary metabolite clothianidin (Blacquière et al. 2012). Based on previous studies indicating negative sublethal effects on worker larvae reared under neonicotinoid-exposure (López et al. 2017), we hypothesized that significantly increased levels of fore wing abnormalities and fluctuating wing asymmetry would be observed in both exposed workers and drones. Furthermore, we expected an increased degree of susceptibility in haploid drones when compared to diploid workers.

## 2. Material and methods

### 2.1. Experimental set-up

The study was performed at the Hasli Ethological Station, Switzerland (46°58'03.359"N 7°23'54.703"E; WGS84 system) using 13 local queenright honey bee colonies established in April 2015 using the shook swarm method (Delaplane et al. 2013). Each colony was initially equipped with five frames containing organic Dadant-sized worker wax foundation (Supplementary data Fig. S1), one laying sister queen, and ~1.8 kg of adult workers (Sandrock et al. 2014a; Williams et al. 2015).

### 2.2. Neonicotinoid exposure

In early May 2015, colonies were randomly allocated to one of two treatments – neonicotinoid (N=6) or control (N=7). According to established methods of in-hive insecticide exposure (Williams et al. 2015; Straub et al. 2016; Forfert et al. 2017), each colony was provided daily with 100 g pollen paste (60% fresh honey bee corbicular pollen, 10% organic honey, and 30% powder sugar) for 50 days to ensure that colonies were exposed for at least two complete brood cycles (Sandrock et al. 2014a; Straub et al. 2016). Pollen patties fed to colonies belonging to the neonicotinoid treatment additionally contained 4.9 ppb thiamethoxam and 2.1 ppb clothianidin (both Sigma-Aldrich), which represent field-realistic concentrations found in agricultural crops and neighboring wild plants (David et al. 2016; Tosi et al. 2018). Concentrations were confirmed by the French National Center for Scientific Research using ultra-high-performance liquid chromatography-tandem mass spectrometry (UHPLC-MS/MS). Throughout the entire exposure period, each colony was fitted with a hive entrance pollen trap to promote pollen paste feeding by reducing forager-collected corbicular pollen (Human et al. 2013).

### 2.3. Source of drones and workers

To ensure that experimental drones were reared during treatment exposure, an additional frame containing organic drone cell wax foundation was added to each colony three weeks after initiation of pollen patty feeding (Rangel and Tarpy, 2016). Eight days later, once drone-sized wax cells were drawn, each queen was caged to its colony's respective drone frame for 48 h to obtain sufficient quantities of individuals of a known age cohort. Immediately after the 48 h period, queens were placed on a worker brood frame within their colony using the previously described procedure to obtain equivalent worker cohorts. Each experimental drone and worker brood frame remained in its respective colony until it was transferred 24 h prior to natural emergence to a laboratory incubator maintained at 34.5 °C and 60% RH in complete darkness (Williams et al. 2013). To ensure that drones would successfully emerge and feed, 50 adult worker attendants from each colony were added to their respective drone frame (Currie, 1987). Post-emergence, both experimental drones and workers remained on their respective brood frames for ~12 h to ensure successful wing

ventilation and maturation (Dickinson, 1999). To ensure wing integrity, drones and workers were swiftly killed by carefully placing individuals into falcon tubes containing cotton soaked in ethyl acetate (Human et al. 2013). Falcon tubes were stored at  $-24^{\circ}\text{C}$  until the wings of the individuals were examined.

#### 2.4. Dissection and wing preparation

Dissection and preparation of experimental honey bee worker and drone wings were conducted during September 2017 and March 2018. Subsamples consisting of five individuals at a time were dissected and mounted to a cover slide to minimize possible negative effects caused by thawing. In total, 7–21 drones and 16–17 workers per colony were assessed by dissecting each in a Petri dish following Carreck et al. (2013). In brief, the thorax was carefully removed from the head and abdomen using dissection scissors. Then, the thorax was dissected longitudinally, to equally separating the left and right body side. Using a fine dissection scissor, the fore and hind wings were removed from the thorax to ensure that each wing base remained intact and could serve as a landmark for measurements (Klingenberg, 2011). Fore wings were placed in EtOH of increasing concentrations (50, 70, 90%) for 5 min before rinsed in distilled water (Meixner et al. 2013), whereas the hind wings were not included for this study.

Fore wings were then carefully mounted on a prepared cover slide to prevent folds or air bubbles underneath the wings which would lead to distortions in landmark measurements (Klingenberg, 2015). To facilitate digitalizing fore wings, each was positioned at the same height and orientation with the aid of a sliding caliper. Once mounted, 0.1 ml of the fixing agent Euparal (Carl Roth®) was carefully applied to each fore wing. Using a stereomicroscope (Olympus SZX7), possible undetected air bubbles in the solution and under the wings were gently removed using a forceps. Then, a coverslip [ $21 \times 26 \text{ mm}$ ] was placed upon each individual fore wing before placing the slide in an incubator at  $30^{\circ}\text{C}$  for 24 h to dry.

#### 2.5. Wing digitalization

Each individual fore wing (left and right) from 152 drones (94 controls and 58 neonicotinoids) and 212 workers (112 controls and 100 neonicotinoids) were analyzed at the Natural History Museum of Bern, Switzerland. Digital images were taken with a high-resolution camera mounted to a digital photo microscope (Keyence VHX, 2000 digital photo microscope with VH-Z20 R/W zoom lens). TPS photo files were built using the software tpsUtil (Rohlf, 2008).

#### 2.6. Abnormalities in wing venation

Following Smith et al. (1997), all fore wing pairs were analyzed to identify wing venation abnormalities, also known as phenodeviants, such as vein protrusions, incomplete veins, as well as extra or missing veins (Smith et al. 1997; Lopuch and Tofilski, 2016). These wing venation abnormalities are likely caused by genetic and/or environmental influences during the development of a particular trait (Smith et al. 1997; Lopuch and Tofilski, 2016). In our study, we quantified the frequency of individual workers and drones from both treatment groups showing at least one wing venation abnormality. Furthermore, we quantified the total number of wing venation abnormalities present in both left and right wing for all individuals.

#### 2.7. Morphometric measurements

The coordinates of 16 standardized, distinguishable landmarks

located at intersections of the different veins and inter-vein areas were recorded and measured twice using the tpsDig software (Fig. S2) (Rohlf, 2001; Klingenberg, 2015) to obtain a measurement error (ME) (Graham et al. 2010). Landmark coordinates were used to derive wing size and Procrustes coordinates that allowed the shape to be characterized, and the type of asymmetry to be determined (Klingenberg, 2015). For drones, we obtained a vector containing 608 sizes corresponding to 152 individuals  $\times$  2 fore wings (right and left)  $\times$  2 replicates. For workers, we obtained a vector containing 848 sizes corresponding to 212 individuals  $\times$  2 fore wings (right and left)  $\times$  2 replicates. After normalizing all wing sizes, samples were aligned to directly compare their shape by obtaining a matrix of Procrustes coordinates containing the vector multiplied by 2 replicates (by 16 landmarks) and 2 dimensions (x and y).

In brief, the Procrustes approach extracted shape information from coordinate data in order to eliminate reflection (right and left wing) and variation in scale, position, and orientation among individual wings (Klingenberg, 2015). First, reflection was removed by changing the sign of the x co-ordinates of all left wings and thereby flipping them horizontally to their mirror images. In a second step, all landmark configurations were scaled to the same centroid size. It is the centroid point (center of gravity) whose co-ordinates are the averages of the respective coordinates of all the landmarks (Klingenberg, 2015). Thus, the position of each wing was standardized by superimposing centroids of all configurations on the origin (0, 0). The last step eliminated variation in orientation by rotating landmark configurations around their shared centroid to achieve an optimal fit of each wing to the consensus (average) configuration (Bookstein, 1991).

#### 2.8. Statistical analyses

All statistical tests and figures were performed and produced using MorphoJ software (Klingenberg, 2011) and NCSS v. 11.05 (NCSS, 2018).

##### 2.8.1. Wing venation abnormalities

Data were tested for normality by using the Shapiro-Wilk's test. We then applied a logistic regression model to determine significant differences among the frequencies of individuals with wing venation abnormalities from different treatments (control and neonicotinoid insecticide) and ploidy levels (diploid and haploid), wherein treatment and ploidy level were included as fixed terms and colony as a random effect. Total wing venation abnormalities per individual were compared using a Kruskal-Wallis ANOVA and post hoc comparison (Bonferroni's test) to test for differences among treatments and sex.

##### 2.8.2. Measurement error

Measurement error can cause a considerable degree of variation in asymmetry. Therefore, repeated measurements of the same individual fore wing were subject to a two-way mixed-model ANOVA [Side (fixed term)  $\times$  Individual (random effect)] with repeated measurements of each side (Ondo et al. 2011). The fixed factor 'Side' tested for significant positive or negative differences between the left and right wing side of an individual regardless of its treatment group (i.e. directional asymmetry) (Graham et al. 2010). Significant effects of the random factor 'Individual' tested for differences in size or shape among individual drones and workers. Directional asymmetry in this context means a greater development of the wing on one side of the plane when compared to the other side (Klingenberg, 2015). It can be quantified by calculating the size and shape difference between averages for left and right wings (Klingenberg, 2015). The significance of the interaction term

'Individual x Side' tested for differences between sides among individuals (i.e. fluctuating asymmetry). Measurement error was considered negligible if the interaction term was significant ( $P < 0.05$ ) and the mean square value for measurement error was substantially smaller than all other mean square values (Graham et al. 2010; Lopuch and Tofilski, 2016).

### 2.8.3. Wing size and wing size asymmetry

Fore wing size was evaluated using centroid size [ $\text{mm}^2$ ] (Dryden and Mardia, 1989). This was calculated as the square root of the sum of the squared distances of the 16 landmarks from the centroid point (Klingenberg et al. 2001). Individual drone and worker fore wing size were defined as the average of the two replicates measured for left and right fore wings; this resulted in one wing size per side for each specimen. Wing size asymmetry was then measured by calculating the difference between the sizes of the right and left fore wing (right-left (R-L)) (Palmer and Strobeck, 2003). Mean value of wing size asymmetry obtained for each individual treatment group represented directional asymmetry, whereas, fluctuating asymmetry of wing size was calculated by subtracting directional asymmetry values from R-L values previously described (Klingenberg, 2015). Colony (random effect) and treatment (fixed term) effects were examined using MANOVAs, whereas, differences in variation were established by using the F-test.

### 2.8.4. Wing shape and wing shape asymmetry

Analysis of fore wing shape was performed by aligning the configurations of landmarks using Procrustes superimposition (Bookstein, 1991). Differences in wing shape between the treatment groups were examined using multivariate factorial ANOVA, with average Procrustes coordinates as random variables (individuals and colonies) and groups as fixed factors (treatments) (Nouvellet et al. 2011).

Because fluctuating asymmetry is the result of random irregularities during development, the magnitude of the shape asymmetry can be used as an estimate of developmental instability (Klingenberg, 2015). Therefore, the Procrustes distance between left and right sides is a straightforward choice for measuring overall wing shape asymmetry. In the presence of directional asymmetry, which is a widespread phenomenon in animals (Pélabon and Hansen, 2008), the Procrustes distance between the left and right sides is the combined result of fluctuating asymmetry and directional asymmetry. Because Procrustes distance is used to determine absolute shape differences, Mahalanobis distance, which compares levels of fluctuating asymmetry between groups, was also employed (Klingenberg, 2015). Therefore, the Procrustes distance was used as a measure to determine absolute shape differences, whereas the Mahalanobis distance measured differences between groups (i.e. sex and treatment groups) relative to the within-group variation (i.e. colony) (Klingenberg, 2011; Lopuch and Tofilski, 2016). Both Procrustes and Mahalanobis distances were calculated by using a canonical variate analysis (CVA), which maximizes the separation of specified groups and enables visualization of variation within treatments and sex (Klingenberg, 2011). Procrustes and Mahalanobis distances, with respective significance ( $P$ ) values for these distance, were generated using permutation tests (10,000 iterations per comparison); a scatter plot of the CVA scores was produced to visualize variation among treatments and sex (Benítez, 2013).

## 3. Results

Summary statistics for all measured variables (i.e. the frequency

of wing venation abnormalities, total wing venation abnormalities, centroid size [mean in  $\text{mm}^2$ ], fluctuating asymmetry of wing size [log] and shape) in both worker and drone fore wings are provided in Table S1.

### 3.1. Wing venation abnormalities

#### 3.1.1. Workers


No significant treatment effect was found between control ( $23.21 \pm 5.05\%$ ) and neonicotinoid-exposed ( $23.92 \pm 4.30\%$ ) workers for frequency of wing venation abnormalities (logistic regression,  $R_2 = 0.008$ ,  $\text{df} = 2$ ,  $Z = -0.86$ ,  $P = 0.39$ ) (mean  $\pm$  standard deviation % (SD); Fig. 1). Likewise, total wing venation abnormalities found in control ( $1 \pm 1-2$ ) and neonicotinoid-exposed ( $1 \pm 1-2$ ) workers did not significantly differ ( $F_{1,49} < 0.01$ ,  $P = 0.93$ ) (median  $\pm$  95% CI, Fig. S3).

#### 3.1.2. Drones

A significant difference was found between the frequency of wing venation abnormalities in control ( $44.53 \pm 16.03\%$ ) and neonicotinoid-exposed ( $70.35 \pm 9.09\%$ ) drones (logistic regression,  $R_2 = 1.15$ ,  $\text{df} = 3$ ,  $Z = 4.85$ ,  $P < 0.001$ , Fig. 1) (mean  $\pm$  % SD). In contrast, no significant difference was observed for the total wing venation abnormalities found between control ( $2 \pm 1-4$ ) and neonicotinoid-exposed ( $2 \pm 14$ ) drones ( $F_{1,79} < 0.2$ ,  $P = 0.66$ ) (median  $\pm$  95% CI, Fig. S3).

#### 3.1.3. Colony-level

Irrespective of the treatment group, a significant difference of individuals showing wing venation abnormalities (logistic



**Fig. 1. Effects of neonicotinoid insecticides on honey bee, *Apis mellifera* (L.), worker and drone fore wing venation.** For each of the four treatment groups, the frequency of wing venation abnormalities was measured. The bar charts show the overall observed frequency of individuals with at least one wing venation abnormality in either left or right wing as well as the mean  $\pm$  standard error. A significant difference (logistic regression,  $P < 0.05$ ) between treatments and sex is indicated by different letters (A, B, C).



regression,  $R_2 = 0.079$ ,  $df = 13$ ,  $Z = -2.42$ ,  $P < 0.015$ ), as well as total wing venation abnormalities ( $F_{12,351} = 5.28$ ,  $P = 0.009$ ), was observed amongst the 13 colonies.

### 3.2. Measurement error

The two-way mixed-model ANOVA with repeated measures of each side in drones and workers showed that mean square values were always substantially smaller than the interaction term, and therefore highly significant (side  $\times$  individual interaction;  $P < 0.001$ , Table 1).

### 3.3. Wing size and wing size asymmetry

#### 3.3.1. Workers

Neonicotinoid-exposure revealed no significant effect on worker wing size or wing size asymmetry ( $F_1 = 3.07$ ,  $P = 0.42$ , Fig. 2A, C; Table 2). Wing size showed significant individual ( $F_{198} = 28.11$ ,  $P < 0.001$ ; Table 2) and colony-level variation ( $F_{12} = 4.43$ ,  $P < 0.001$ ; Table 2), with worker fore wings having a mean centroid size of 651.24 mm<sup>2</sup>. Significant directional asymmetry of wing size was observed ( $F_1 = 4.98$ ,  $P < 0.03$ ; Tables 2 and 3), with the right wings in both treatment groups on average ~0.014% larger than the left wings. Regardless of treatment group, significant fluctuating asymmetry of wing size was observed for all individuals ( $F_{211} = 57.14$ ,  $P < 0.001$ ; Tables 2 and 3).

#### 3.3.2. Drones

Similar to workers, no significant treatment effect was observed for either wing size or wing size asymmetry ( $F_1 = 0.48$ ,  $P = 0.49$ , Fig. 2B, D; Table 2). Significant individual ( $F_{150} = 29.97$ ,  $P < 0.001$ , Table 2) and colony-level variation ( $F_9 = 14.94$ ,  $P < 0.001$ ; Table 2) were also observed for drone wing size; mean centroid size was 887.80 mm<sup>2</sup>. Significant directional asymmetry of wing size was detected in drones ( $F_1 = 15$ ,  $P < 0.002$ ; Tables 2 and 3), with the right fore wing in both treatment groups ~0.067% larger than the left one. Additionally, significant fluctuating asymmetry of wing size was observed in all experimental drones ( $F_{152} = 69.94$ ,  $P < 0.001$ ; Tables 2 and 3).

**Table 1**  
Calculated measurement errors using Procrustes ANOVA of centroid size and shape of honey bee, *Apis mellifera* (L.), drone and worker fore wings. Results obtained from the Procrustes analysis were used to assess measurement error on the fore wing samples that had been measured twice. The analyses evaluated five effects for both centroid size and shape: 1. 'Sex' - female workers or male drones, 2. 'Individual' - tested for differences among individuals ( $N = 364$ ), 3. 'Side' - tested for directional asymmetry within individuals (i.e. one wing side consistently different from the other), 4. 'Individual  $\times$  Side' - tested for differences in fluctuating asymmetry among individuals, and 5. Measurement error was not of concern as the 'Individual  $\times$  Side' interaction was highly significant and the corresponding mean square substantially exceeded measurement error.

Fore wing				
	Mean Square (MS)	df	F	P (param.)
<b>Centroid size [MS in mm<sup>2</sup>]</b>				
Sex	2670218.88	1	2448.64	<0.0001
Individual	1090.49	362	31.70	<0.0001
Side	602.30	1	17.51	<0.0001
Individual $\times$ Side	34.40	363	62.92	<0.0001
Measurement Error	0.55	727		
<b>Shape [Procrustes MS]</b>				
Sex	0.0139904	28	193.15	<0.0001
Individual	0.0000724	10136	11.88	<0.0001
Side	0.0000519	28	8.51	<0.0001
Individual $\times$ Side	0.0000061	10164	17.19	<0.0001
Measurement Error	0.0000004	20356		

### 3.4. Wing shape and wing shape asymmetry

#### 3.4.1. Workers


In contrast to wing size, a significant treatment effect on wing shape was observed ( $F_{28} = 2.54$ ,  $P < 0.001$ ; Fig. 2E, Table 2). Fore wing shape among all individuals ( $F_{554} = 13.13$ ,  $P < 0.001$ ; Table 2) and colonies ( $F_{336} = 4.54$ ,  $P < 0.001$ ; Table 2) significantly differed. Significant directional asymmetry of wing shape was observed for all individual workers ( $F_{28} = 10.8$ ,  $P < 0.001$ ; Table 2). Regardless of treatment group, significant fluctuating asymmetry of wing size was observed for all individuals ( $F_{5908} = 18.07$ ,  $P < 0.001$ ; Table 2). A highly significant Procrustes distance of 0.0039 was revealed ( $P < 0.001$ ; Table 4), with a fluctuating asymmetry of wing shape between left and right wings being larger in neonicotinoid-exposed workers (0.0028) than in controls (0.0030) (Procrustes distance; Fig. 3; Table 4). Likewise, a significant difference was also observed for the Mahalanobis distance measured between treatment groups ( $P < 0.001$ ; Table 4), which was 1.063. Control and neonicotinoid-exposed workers showed significant fluctuating asymmetry of wing shape between left and right wings of 0.666 and 0.721 respectively, which indicates increased shape asymmetry in the neonicotinoid-exposed workers (Mahalanobis distances,  $P < 0.001$ ; Table 4).

#### 3.4.2. Drones

Similar to workers, a significant effect of neonicotinoids on wing shape was observed ( $F_{28} = 6.45$ ,  $P < 0.001$ , Fig. 2F; Table 2). Again, fore wing shape among all individuals ( $F_{4200} = 13.13$ ,  $P < 0.001$ ; Table 2) and colonies ( $F_{252} = 7.67$ ,  $P < 0.001$ ; Table 2) significantly differed. Furthermore, a significant difference was observed for directional asymmetry of wing shape among individuals ( $F_{28} = 1.95$ ,  $P < 0.001$ ; Table 2). Significant fluctuating asymmetry of wing shape was also observed among experimental drones ( $F_{4228} = 16.25$ ,  $P < 0.001$ ; Table 2). A significant Procrustes distance of 0.0092 was revealed ( $P < 0.001$ ; Table 4), with the fluctuating asymmetry of wing shape between left and right wings larger in neonicotinoid drones (0.0023) than in controls (0.0016) (Procrustes distance, Fig. 3; Table 4). The Mahalanobis distance measured between treatment groups was 2.0247 and was significant ( $P < 0.001$ ; Table 4). Control and neonicotinoid-exposed drones showed fluctuating asymmetry of wing shape between left and right fore wings of 0.409 and 0.599, respectively, which indicates increased shape asymmetry in neonicotinoid-exposed individuals (Mahalanobis distances;  $P < 0.001$ , Table 4).

### 3.5. Comparison between workers and drones

Drones had a significantly higher frequency of wing venation abnormalities compared to workers (logistic regression;  $R_2 = 0.56$ ,  $df = 2$ ,  $Z = 2.2$ ,  $P < 0.03$ ; Fig. 1). In addition, drones experienced significantly more total abnormalities when compared to workers ( $F_{3,128} = 6.68$ ,  $P < 0.001$ , Fig. S3). A significant difference was observed between worker and drone wing size (One way ANOVA;  $P < 0.001$ ), with mean centroid wing size being  $650.48 \pm 11.48$  mm<sup>2</sup> and  $888.53 \pm 17.15$  mm<sup>2</sup> for the workers and drones, respectively (mean  $\pm$  SD; Table S1). Therefore, the average drone fore wing was 26.8% larger than that of a worker's. Based on Procrustes distances, the CVA scores revealed significant differences in fore wing shape between workers and drones as well as between treatments ( $P < 0.001$ ), which resulted in two non-overlapping clouds of points clearly discriminating workers from drones (Fig. 3). The sum of the first (97.04%) and second (2.45%) CV account for over (99.49%) of the relative between-groups variation and accordingly, it was sufficient to examine only the computed plot of the first two CVs. There was a significant difference for both the Procrustes and



**Fig. 2.** Effects of neonicotinoid insecticides on honey bee, *Apis mellifera*, worker (A, C, E) and drone (B, D, F) fore wing size and shape. All variables were measured using 16 landmarks from each fore wing. Wing size was evaluated using the centroid size [mm<sup>2</sup>], A & B) whereas fluctuating asymmetry of wing size (centroid size [log], C & D) was then measured by calculating the difference between the sizes of the right and left fore wings (R–L). No significant treatment effect was observed either workers or drones for either of these parameters ( $P > 0.05$ ). Analysis of fore wing shape was performed by aligning the configurations of landmarks using Procrustes superimposition (Procrustes distance (P.D.) in E & F). Significant differences were observed for both workers and drones ( $P < 0.001$ ). The boxplots show the inter-quartile-range (box), the median (black line within box), and outliers (dots). A significant difference ( $P < 0.001$ ) between treatments is indicated by \*\*\*.

Mahalanobis distance between workers and drones ( $P < 0.001$ ; Table 4), with the determined distances being 0.033 and 9.33, respectively. In addition, our results revealed that the degree of

fluctuating asymmetry of wing shape was significantly higher in drones (0.0092) than in workers (0.0039) (Procrustes Distance,  $P < 0.001$ ; Table 4).

**Table 2**

**Summary of honey bee drone and worker, *Apis mellifera* (L.), fore wing size and shape for control and neonicotinoid treatment groups.** Degrees of freedom (df), the sum of squares, mean squares, and *F* and *P*-values are presented for random effects 'Individual' and 'Colony', as well as for fixed effects 'Treatment' (control and neonicotinoid) and 'Side' (e.g. right and left fore wing). The interaction term was included to indicate the level of fluctuating asymmetry. Treatment effects were observed for shape ( $P < 0.001$ ) but not size ( $P = 0.4190$  for workers,  $P = 0.4906$  for drones).

Sex	Source	Sum of Square (SS)	quare (MS)	df	<i>F</i>	<i>P</i>
<b>Centroid Size</b>						
<b>Worker</b>	Treatment	2640.75	264.75	1	3.07	0.4190
	Colony	45772.44	3814.37	12	4.43	<0.0001
	Individual	170406.12	86.64	198	28.11	<0.0001
	Side	152.48	152.48	1	4.98	0.0267
	Individual x Side	6460.86	30.62	211	57.14	<0.0001
<b>Drone</b>	Treatment	305.38	305.38	1	0.48	0.4906
	Colony	85947.93	9549.77	9	14.94	<0.0001
	Individual	176079.34	1173.86	150	29.97	<0.0001
	Side	587.68	587.68	1	15.00	0.0020
	Individual x Side	5914.91	39.17	151	69.94	<0.0001
<b>Shape</b>						
<b>Worker</b>	Treatment	0.003231	0.000115	28	2.54	<0.0001
	Colony	0.069355	0.000206	336	4.54	<0.0001
	Individual	0.252322	0.000046	554	8.67	<0.0001
	Side	0.001588	0.000057	28	10.8	<0.0001
	Individual x Side	0.031028	0.000005	5908	18.07	<0.0001
<b>Drone</b>	Treatment	0.012217	0.000436	28	6.45	<0.0001
	Colony	0.130821	0.000519	252	7.67	<0.0001
	Individual	0.000095	0.000095	4200	13.13	<0.0001
	Side	0.000393	0.000014	28	1.95	0.0024
	Individual x Side	0.000007	0.000007	4228	16.25	<0.0001

**Table 3**


**Summary of obtained sizes for honey bee, *Apis mellifera* (L.), drone and worker fore wings, and mean asymmetry of fore wing size.** Centroid size represented fore wing size and was calculated by the square root of the sum of the squared distances of the 16 landmarks from the centroid (center of gravity). Results revealed that the right wing was larger than the left wing for both treatment groups, which resulted in a significant degree of fluctuating asymmetry of wing size for both controls and neonicotinoids ( $P < 0.001$ ).

Sex	Treatment	Wing Side	Centroid Size (log)			Wing Size Asymmetry (log)		
			Mean	Standard Error	<i>P</i>	Mean	Standard Error	<i>P</i>
<b>Worker</b>	Control	Right	2.8139	1.0588	0.0316	0.0022	0.0019	<0.001
		Left	2.8136	1.0630				
	Neonicotinoid	Right	2.8125	0.9966	0.0081	0.0014	0.0010	<0.001
		Left	2.8120	0.9760				
<b>Drone</b>	Control	Right	2.9488	1.2455	0.0107	0.0024	0.0027	<0.001
		Left	2.9478	1.2677				
	Neonicotinoid	Right	2.9495	1.2125	0.0016	0.0021	0.0015	<0.001
		Left	2.9485	1.2113				

#### 4. Discussion

Our data show for the first time higher developmental instability of male haploid insects to neonicotinoids compared to diploid females. This provides empirical support to the haploid susceptibility hypothesis that points to increased susceptibility of haploids to environmental stressors (O'Donnell and Beshers, 2004). Our data also provide general evidence that field-realistic concentrations of two widely employed neonicotinoid insecticides can negatively affect developmental stability in honey bees. Thiamethoxam and clothianidin significantly increased the frequency of wing venation anomalies, as well as the degree of fluctuating asymmetry in fore wings. This suggests that neonicotinoid insecticides may interfere with essential genetic cascades responsible for regulating developmental stability during larval insect development (De Celis and Diaz-Benjumea, 2003), thereby contributing to the overwhelming evidence for effects of these chemicals on non-target organisms.

There is a consensus that abnormalities in wing venation patterns of insects are caused by both genetic and environmental factors (Mazeed, 2011; Lopuch and Tofilski, 2016). Regardless of the factor, the mechanisms explaining such wing venation abnormalities in holometabolous insects must occur during metamorphosis, when the rudiments of venation are still plastic (Truman and Riddiford, 2019). Minor deviations or additional fragments can be interpreted as atavistic phenomena (Zanni and Opitz, 2013), which were observed in our study for both honey bee drones and workers. Since control drones were more likely to show abnormalities compared to workers in the absence of any experimental environmental stressor, our results provide empirical support that recessive alleles can explain the higher frequency of such abnormalities (Bährmann, 1963). This also suggests that minor wing abnormalities are unlikely to interfere with flight performance. Indeed, about 20% of workers had abnormalities in completely functional colonies, irrespective of neonicotinoid treatment.



Interestingly, our data revealed that the fluctuating wing shape asymmetry was significantly higher in drones when compared to workers. Furthermore, the frequency of workers showing wing venation abnormalities, as well as the number of abnormalities present per individual, was not significantly influenced by neonicotinoid exposure. In sharp contrast, neonicotinoid exposure significantly increased the frequency, but not the total number of wing venation abnormalities in drones. Similar results for wing abnormalities and mean number of abnormalities have been reported for honey bees (Lopuch and Tofilski, 2016). However, for the first time, we revealed an increased frequency of wing abnormalities and fluctuating wing asymmetry for haploid neonicotinoid-exposed drones when compared to diploid workers. This may be due to lack of heterozygosity at loci relevant during metamorphosis. Indeed, the neonicotinoids may interfere with genetic pathways relevant for the development of wing veins (Bier, 2000), because neonicotinoid-exposed bees also have to activate the detoxification pathways (Claudianos et al. 2006), such as cytochrome P450s or glutathione S-transferases (Li et al. 2007).

**Table 4**  
**Summary and comparison of Procrustes distances measured for honey bee, *Apis mellifera* (L.), worker and drone fore wings in control and neonicotinoid treatment groups.** The degree of fluctuating shape asymmetry between left and right fore wing of each treatment was measured using Procrustes distances. Neonicotinoid-exposed workers and drones revealed a significantly higher degree of Procrustes distances ( $P < 0.001$ ).

Worker	Procrustes Distances		P-values for Procrustes Distances		Mahaalanobis Distances		P -values for Mahalanobis Distances			
	Control Left	Control Right	Neonicotinoid Left	Control Left	Control Right	Neonicotinoid Left	Control Left	Control Right	Neonicotinoid Left	Control Right
Sex	0.033			<0.001		9.3268				<0.001
Treatment	0.0039			<0.001		1.063				
Drone	Treatment and Side		Neonicotinoid Left	Control Left	Control Right	Neonicotinoid left	Control Left	Control Right	Neonicotinoid Left	Control Right
	Control Right	0.0028		0.02		0.666			<0.0001	
	Neonicotinoid Left	0.0041		<0.0001		1.1055		1.1855	<0.0001	
	Neonicotinoid Right	0.0051	0.003	<0.0001	<0.0001	1.3089		1.0645	<0.0001	<0.0001
	Treatment	0.0092		<0.001		2.0257			<0.001	
	Treatment and Side		Neonicotinoid Left	Control Left	Control Right	Neonicotinoid left	Control Left	Control Right	Neonicotinoid Left	Control Right
	Control Right	0.0016		0.99		0.4089			0.99	
	Neonicotinoid Left	0.0092		<0.0001	<0.0001	2.2695		2.38 27	<0.0001	<0.0001
	Neonicotinoid Right	0.0088	0.0023	<0.0001	<0.0001	2.1859		2.2318	0.0004	0.9981
	Treatment	0.0093		<0.0001	<0.0001	0.97		0.5991	<0.0001	<0.0001



The need for activation of additional gene cascades in exposed individuals may limit resources, which could otherwise be allocated to the core pathways essential for metamorphosis (Belles and Santos, 2014; Truman and Riddiford, 2019). As wing shape and wing venation patterns are governed by independent molecular mechanisms (Shimmi et al. 2014), our data suggest that neonicotinoid insecticides may negatively affect developmental homeostasis by interfering with at least two different mechanisms.

Even though neonicotinoid-exposed haploid drones were more susceptible, the observed degree of wing malformations in both controls and treatments is unlikely to interfere with flight abilities. Thus, the reported impaired flight ability of neonicotinoid-exposed bees (Blanken et al. 2015; Tison et al. 2016; Tosi et al. 2017) is more likely to result from an impact on behavior. This appears very plausible because neonicotinoids are neurotoxins in the first place (Blacqui re et al. 2012).

Our data show that fore wing size and shape, as well as wing venation abnormalities, can significantly vary among colonies, regardless of treatment exposure. Considering that all colonies were maintained under the same conditions (common garden approach), it appears likely that genetic variation may be responsible. Such variation in sensitivity towards insecticides is known (Miyo et al. 2000). Yet, our results provide further evidence that the detoxification capacities in honey bee colonies can significantly vary (Sandrock et al. 2014b) and subsequently highlight the importance of considering both individual and colony-level genetics when interpreting data of toxicological studies. Moreover, the data confirm a significant level of directional asymmetry of wing size and shape, in favor of larger right wings. Differences in left and right body sides have already been observed in several haploid-diploid insects (Klingenberg, 2019), and is known to be widespread throughout the animal kingdom (Palmer, 1996). These differences are most likely explained by the existence of a left-right axis that reflects distinct positional identities of the wing imaginal discs on either body side (Klingenberg et al. 1998); however, the overall small magnitude of directional asymmetry precludes directional asymmetry from having any major adaptive role or consequences (P labon and Hansen, 2008).

In conclusion, our data show that neonicotinoids can interfere with honey bee metamorphosis and that haploids are more susceptible. This creates demand for a better understanding of the possible impact of these insecticides on other factors during metamorphosis (e.g. spermatogenesis (Straub et al. 2016)). Further, haplodiploid insect species and genetically less diverse populations may be more at risk, which seems relevant for conservation efforts (Reed and Frankham, 2003; Winfree, 2010). Future risk assessments should, therefore, consider the genetic basis of susceptibility to insecticides and the weakest link within a species.

## Funding

Financial support was provided by the Bundesamt f r Umwelt (BAFU) to L.S., G.R.W. and P.N., the Vinetum Foundation to P.N., the Swiss National Science Foundation (SNSF) project 31003A\_169751 to G.R.W., the Foundation for Food and Agriculture Research Pollinator Health Fund grant 549003 to G.R.W., and the USDA National Institute of Food and Agriculture Multi-state Hatch project 1014681 to G.R.W.

## Data accessibility

The complete raw data can be found on the Dryad repository (<https://doi.org/10.5061/dryad.7d7wm37r0>).

## Authors' contributions

L.S., G.R.W. & P.N. conceived the study; A.F., S.B., G.R.W. & L.S. carried out the experiment and collected data; P.N. provided materials; A.F. & L.S. designed the statistical analysis and wrote the statistical methods and analyzed data; All authors contributed to the interpretation of results and writing of the manuscript.

## Declaration of competing interest

We declare no competing interests.

## Acknowledgments

We sincerely thank Hannes Baur from the Natural History Museum of Bern, Switzerland, for his kind support throughout the entire project and for his valuable input regarding methods, analyses, and interpretation of results.

## Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.chemosphere.2019.125145>.

## References

- Abbo, P.M.P.M., Kawasaki, J.K., Hamilton, M., Cook, S.C., Degrandi-hoffman, G., Li, W.F.L., et al., 2016. Effects of Imidacloprid and *Varroa destructor* on survival and health of European honey bees, *Apis mellifera*. *Insect Sci.* 00, 1–11.
- Baer, B., Schmid-Hempel, P., 2003. Bumblebee workers from different sire groups vary in susceptibility to parasite infection. *Ecol. Lett.* 6, 106–110.
- B hrmann, R., 1963. Zum Vorkommen von Anomalien im Fl gelge der der Honigbiene. *Arch. Bienenkd* 40, 49–58.
- Beasley, D.A.E., Bonisoli-Alquati, A., Mousseau, T.A., 2013. The use of fluctuating asymmetry as a measure of environmentally induced developmental instability: a meta-analysis. *Ecol. Indic.* 30, 218–226.
- Belles, X., Santos, C.G., 2014. The MEKRE93 (Methoprene tolerant-Kr ppel homolog 1-E93) pathway in the regulation of insect metamorphosis, and the homology of the pupal stage. *Insect Biochem. Mol. Biol.* 52, 60–68.
- Ben tez, H.A., 2013. Assessment of patterns of fluctuating asymmetry and sexual dimorphism in carabid body shape. *Neotrop. Entomol.* 42, 164–169.
- Bier, E., 2000. Drawing lines in the *Drosophila* wing: Initiation of wing vein development. *Current Opinion in Genetics and Development* 10 (4), 393–398.
- Blacqui re, T., Smagghe, G., van Gestel, C. A. M., Mommaerts, V., 2012. Neonicotinoids in bees: a review on concentrations, side-effects and risk assessment. *Ecotoxicology* 21, 973–992.
- Blanken, L.J., Langevelde, F. Van, Dooremalen, C. Van, 2015. Interaction between *Varroa destructor* and imidacloprid reduces flight capacity of honeybees. *Proc. R. Soc. B* 282, 1–9.
- Bookstein, F.L., 1991. *Morphometric Tools for Landmark Data*. Cambridge University Press.
- Calderone, N.W., 2012. Insect pollinated crops, insect pollinators and US agriculture: trend analysis of aggregate data for the period 1992–2009. *PLoS One* 7, e37235.
- Carreck, N.L., Andree, M., Brent, C.S., Cox-foster, D., Dade, H.A., Ellis, J.D., et al., 2013. Standard methods for *Apis mellifera* anatomy and dissection. *J. Apic. Res.* 52, 1–40.
- Carreck, N.L., Ratnieks, F.L.W., 2014. The dose makes the poison: have “field realistic” rates of exposure of bees to neonicotinoid insecticides been overestimated in laboratory studies? *J. Apic. Res.* 53, 607–614.
- De Celis, J.F., Diaz-Benjumea, F.J., 2003. Developmental basis for vein pattern variations in insect wings. *Int. J. Dev. Biol.* 47, 653–663.
- Chang, X., Zhai, B., Wang, M., Wang, B., 2007. Relationship between exposure to an insecticide and fluctuating asymmetry in a damselfly (Odonata, Coenagrionidae). *Hydrobiologia* 586, 213–220.
- Claudianos, C., Biswas, S., Schuler, M.A., Berenbaum, M.R., Feyereisen, R., Oakeshott, J.G., 2006. A deficit of detoxification enzymes: pesticide sensitivity and environmental response in the honeybee. *Insect Mol. Biol.* 15, 615–636.
- Combes, S.A., Daniel, T.L., 2003. Flexural stiffness in insect wings I. Scaling and the influence of wing venation. *J. Exp. Biol.* 206, 2979–2987.
- Currie, R.W., 1987. The biology and behaviour of drones. *Bee World* 68, 129–143.
- David, A., Bot as, C., Abdul-Sada, A., Nicholls, E., Rotheray, E.L., Hill, E.M., et al., 2016. Widespread contamination of wildflower and bee-collected pollen with complex mixtures of neonicotinoids and fungicides commonly applied to crops. *Environ. Int.* 88, 169–178.
- Delaplane, K.S., Steen, J. Van Der, Guzman-Novoa, E., 2013. Standard methods for estimating strength parameters of *Apis mellifera* colonies. *J. Apic. Res.* 52, 1–12.
- Derecka, K., Blythe, M.J., Malla, S., Genereux, D.P., Guffanti, A., Pavan, P., et al., 2013.

- Transient exposure to low levels of insecticide affects metabolic networks of honeybee larvae. *PLoS One* 8 (7), e68191.
- Dickinson, M.H., 1999. Insect flight. *Science* 284 (5422), 1954–1960.
- Dryden, I.L., Mardia, K.V., 1989. Statistical analysis of shape. *Biometrika* 76, 271–281.
- EFSA, 2014. EFSA Guidance Document on the risk assessment of plant protection products on bees (*Apis mellifera*, *Bombus* spp. and solitary bees). *EFSA J* 11, 268.
- Evans, J.D., Shearman, D.C.A., Oldroyd, B.P., 2004. Molecular basis of sex determination in haplodiploids. *Trends Ecol. Evol.* 19, 1–3.
- Forfert, N., Troxler, A., Retschnig, G., Gauthier, L., Straub, L., Moritz, R.F.A.R.F.A., et al., 2017. Neonicotinoid pesticides can reduce honeybee colony genetic diversity. *PLoS One* 12, 1–14.
- Garibaldi, L.A., Carvalheiro, L.G., Vaissière, B.E., Gemmill-herren, B., Hipólito, J., Freitas, B.M., et al., 2016. Mutually beneficial pollinator diversity and crop yield outcomes in small and large farms. *Science* 351 (6271), 388–391.
- Gerard, M., Michez, D., Debat, V., Fullgrabe, L., Meeus, I., Piot, N., et al., 2018. Stressful conditions reveal decrease in size, modification of shape but relatively stable asymmetry in bumblebee wings. *Sci. Rep.* 8 (1), 15169.
- Goulson, D., 2013. An overview of the environmental risks posed by neonicotinoid insecticides. *J. Appl. Ecol.* 50, 977–987.
- Graham, J.H., Raz, S., Hel-Or, H., Nevo, E., 2010. Fluctuating asymmetry: methods, theory, and applications. *Symmetry* 2, 466–540.
- Hallmann, C.A., Sorg, M., Jongejans, E., Siepel, H., Hofland, N., Schwan, H., et al., 2017. More than 75 percent decline over 27 years in total flying insect biomass in protected areas. *PLoS One* 12 (10), e0185809.
- Higginson, A.D., Barnard, C.J., 2004. Accumulating wing damage affects foraging decisions in honeybees (*Apis mellifera* L.). *Ecol. Entomol.* 29, 52–59.
- Human, H., Brodschneider, R., Dietemann, V., Dively, G., Ellis, J.D., Forsgren, E., et al., 2013. Miscellaneous standard methods for *Apis mellifera* research. *J. Apic. Res.* 52, 1–56.
- Klingenberg, C.P., 2011. MorphoJ: an integrated software package for geometric morphometrics. *Mol. Ecol. Resour.* 11, 353–357.
- Klingenberg, C.P., 2015. Analyzing fluctuating asymmetry with geometric morphometrics: concepts, methods, and applications. *Symmetry* 7, 843–934.
- Klingenberg, C.P., 2019. Phenotypic plasticity, developmental instability, and robustness: the concepts and how they are connected. *Front. Ecol. Evol.* 7, 1–15.
- Klingenberg, C.P., Badyaev, A.V., Sowry, S.M., Beckwith, N.J., 2001. Inferring developmental modularity from morphological integration: analysis of individual variation and asymmetry in bumblebee wings. *Am. Nat.* 157, 1–13.
- Klingenberg, C.P., McIntyre, G.S., Zaklan, S.D., 1998. Left-right asymmetry of fly wings and the evolution of body axes. *Proc. R. Soc. Biol. Sci.* 265, 1255–1259.
- Klingenberg, C.P., Nijhout, H.F., 1999. Genetics of fluctuating asymmetry: a developmental model of developmental instability. *Evolution* 53, 358–375.
- Li, X., Schuler, M.A., Berenbaum, M.R., 2007. Molecular mechanisms of metabolic resistance to synthetic and natural xenobiotics. *Annu. Rev. Entomol.* 52, 231–253.
- López, J.H., Krainer, S., Engert, A., Schuehly, W., Riessberger-Gallé, U., Crailsheim, K., 2017. Sublethal pesticide doses negatively affect survival and the cellular responses in American foulbrood-infected honeybee larvae. *Sci. Rep.* 7, 40853.
- Lopuch, S., Tofilski, A., 2016. The relationship between asymmetry, size and unusual venation in honey bees (*Apis mellifera*). *Bull. Entomol. Res.* 106, 304–313.
- Mazeed, A.M., 2011. Anomalies and asymmetry of wing venation pattern in Carniolan and Egyptian bee populations in Egypt. *Egypt. J. Agric. Res.* 4, 149–161.
- Meixner, M., Pinto, M.A., Bouga, M., Kryger, P., Ivanova, E., Fuchs, S., 2013. Standard methods for characterising subspecies and ecotypes of *Apis mellifera*. *J. Apic. Res.* 52, 1–28.
- Miyo, T., Akai, S., Oguma, Y., 2000. Seasonal fluctuation in susceptibility to insecticides within natural populations of *Drosophila melanogaster*. Empirical observations of fitness costs of insecticide resistance. *Genes Genet. Syst.* 75, 97–104.
- Møller, A.P., 1997. Developmental stability and fitness: a review. *Am. Nat.* 149, 916–932.
- Müller, T., Prosche, A., Müller, C., 2017. Sublethal insecticide exposure affects reproduction, chemical phenotype as well as offspring development and antennae symmetry of a leaf beetle. *Environ. Pollut.* 230, 709–717.
- NCSS, 2018. NCSS 12 Statistical Software.
- Neumann, P., Carreck, N., 2010. Honey bee colony losses. *J. Apic. Res.* 49 (1), 1–6.
- Neumann, P., Frouz, J., Helenius, J., Sarthou, J., Klein, A., Genersch, E., et al., 2015. *Ecosystem Services, Agriculture and Neonicotinoids*. EASAC Policy Report 26, p. 62. April 2015.– 649 ISBN: 978-3-8047-3437-1.
- Nouvellet, P., Ramirez-Sierra, M.J., Dumonteil, E., Gourbière, S., 2011. Effects of genetic factors and infection status on wing morphology of *Triatoma dimidiata* species complex in the Yucatán peninsula, Mexico. *Infect. Genet. Evol.* 11, 1243–1249.
- O'Donnell, S., Beshers, S.N., 2004. The role of male disease susceptibility in the evolution of haplodiploid insect societies. *Proc. Biol. Sci.* 271, 979–983.
- Ondo, N., Abaga, Z., Alibert, P., Dousset, S., Savadogo, P.W., Savadogo, M., et al., 2011. Insecticide residues in cotton soils of Burkina Faso and effects of insecticides on fluctuating asymmetry in honey bees (*Apis mellifera* Linnaeus). *Chemosphere* 83, 585–592.
- Palmer, A.R., 1996. From symmetry to asymmetry: phylogenetic patterns of asymmetry. *Proc. Natl. Acad. Sci.* 93, 14279–14286.
- Palmer, A.R., Strobeck, C., 2003. Fluctuating asymmetry analyses revisited. In: Polak, M. (Ed.), *Developmental Instability: Causes and Consequences*. Oxford University Press, New York, NY, USA, pp. 279–319.
- Pélabon, C., Hansen, T.F., 2008. On the adaptive accuracy of directional asymmetry in insect wing size. *Evolution* 62 (11), 2855–2867.
- Potts, S.G., Imperatriz-Fonseca, V., Ngo, H.T., Aizen, M.A., Biesmeijer, J.C., Breeze, T.D., et al., 2016. Safeguarding pollinators and their values to human well-being. *Nature* 540 (7632), 220–229.
- Prado-Silva, A., Nunes, L.A., dos Santos, J.M., Affonso, P.R.A., de, M., Waldschmidt, A.M., 2018. Morphogenetic alterations in *Melipona quadrifasciata anthidioides* (Hymenoptera: apidae) associated with pesticides. *Arch. Environ. Contam. Toxicol.* 74, 627–632.
- Rangel, J., Tarpy, D.R., 2016. In-hive miticides and their effect on queen superseding and colony growth in the honey bee (*Apis mellifera*). *J. Environ. Anal. Toxicol.* 6 (3), 1–7.
- Reed, D.H., Frankham, R., 2003. Correlation between fitness and genetic diversity. *Conserv. Biol.* 17 (1), 230–237.
- Retschnig, G., Williams, G.R., Mehmman, M.M., Yañez, O., de Miranda, J.R., Neumann, P., 2014. Sex-specific differences in pathogen susceptibility in honey bees (*Apis mellifera*). *PLoS One* 9 (1), e85261.
- Rohlf, F., 2001. TPSDig: Digitize Landmarks from Image Files, Scanner, or Video.
- Rohlf, F., 2008. tpsUtil and tpsDig [cd rom].
- Rosa, A. de S., Teixeira, J.S.G., Vollet-Neto, A., Queiroz, E.P., Blochtein, B., Pires, C.S.S.S., et al., 2016. Consumption of the neonicotinoid thiamethoxam during the larval stage affects the survival and development of the stingless bee. *Scaptotrigona aff. depilis*. *Apidologie* 47, 729–738.
- Ross, L., Davies, N.G., Gardner, A., 2019. How to make a haploid male. *Evol. Lett.* 3, 173–184.
- Sandrock, C., Tanadini, L.G., Pettis, J.S., Biesmeijer, J.C., Potts, S.G., Neumann, P., 2014a. Sublethal neonicotinoid insecticide exposure reduces solitary bee reproductive success. *Agric. For. Entomol.* 16 (2), 119–128.
- Sandrock, C., Tanadini, M., Tanadini, L.G., Fauser-Misslin, A., Potts, S.G., Neumann, P., 2014b. Impact of chronic neonicotinoid exposure on honeybee colony performance and queen superseding. *PLoS One* 9 (8), e103592.
- Schreinemachers, P., Tipraqsa, P., 2012. Agricultural pesticides and land use intensification in high, middle and low income countries. *Food Policy* 37, 616–626.
- Shimmi, O., Matsuda, S., Hatakeyama, M., 2014. Insights into the molecular mechanisms underlying diversified wing venation among insects. *Proc. R. Soc. Biol. Sci.* 281, 1–8.
- Simon-Delso, N., Amaral-Rogers, V., Belzunces, L.P., Bonmatin, J.M., Chagnon, M., Downs, C., et al., 2015. Systemic insecticides (Neonicotinoids and fipronil): trends, uses, mode of action and metabolites. *Environ. Sci. Pollut. Res.* 22, 5–34.
- Smith, D.R., Crespi, B.J., Bookstein, F.L., 1997. Fluctuating asymmetry in the honey bee, *Apis mellifera*: effects of ploidy and hybridization. *J. Evol. Biol.* 10, 551–574.
- Stanley, D.A., Michael, P., Garratt, D., Wickens, J.B., Wickens, V.J., Potts, S.G., et al., 2015. Neonicotinoid pesticide exposure impairs crop pollination services provided by bumblebees. *Nature* 528 (7583), 548–550.
- Straub, L., Villamar-bouza, L., Bruckner, S., Chantawannakul, P., Gauthier, L., Khongphinitbunjong, K., et al., 2016. Neonicotinoid insecticides can serve as inadvertent insect contraceptives. *R. Soc. Proc. B* 283 (1835), 20160506.
- Straub, L., Williams, G.R.G.R., Pettis, J., Fries, I., Neumann, P., 2015. Superorganism resilience: eusociality and susceptibility of ecosystem service providing insects to stressors. *Curr. Opin. Insect Sci.* 12, 109–112.
- Strobl, V., Yañez, O., Straub, L., Albrecht, M., Neumann, P., 2019. Trypanosomatid parasites infecting managed honeybees and wild solitary bees. *Int. J. Parasitol.* 49, 605–613.
- Tison, L., Hahn, M.L., Holtz, S., Rößner, A., Greggers, U., Bischoff, G., et al., 2016. Honey bees' behavior is impaired by chronic exposure to the neonicotinoid thiacloprid in the field. *Environ. Sci. Technol.* 50, 7218–7227.
- Tosi, S., Burgio, G., Nieh, J.C., 2017. A common neonicotinoid pesticide, thiamethoxam, impairs honey bee flight ability. *Sci. Rep.* 7, 1201.
- Tosi, S., Costa, C., Vesco, U., Quaglia, G., Guido, G., 2018. A 3-year survey of Italian honey bee-collected pollen reveals widespread contamination by agricultural pesticides. *Sci. Total Environ.* 615, 208–218.
- Truman, J.W., Riddiford, L.M., 2019. The evolution of insect metamorphosis: a developmental and endocrine view. *Philos. Trans. R. Soc. Biol. Sci.* 374, 20190070.
- Williams, G.R., Alaux, C., Costa, C., Csáki, T., Doublet, V., Eisenhardt, D., et al., 2013. Standard methods for maintaining adult *Apis mellifera* in cages under in vitro laboratory conditions. *J. Apic. Res.* 52, 1–36.
- Williams, G.R., Troxler, A., Retschnig, G., Roth, K., Yañez, O., Shutler, D., et al., 2015. Neonicotinoid pesticides severely affect honey bee queens. *Sci. Rep.* 5, 14621.
- Willmore, K.E., Hallgrímsson, B., 2005. Within individual variation: developmental noise versus developmental stability. *Variation - A Central Concept in Biology*. Elsevier Inc., pp. 191–218.
- Winfree, R., 2010. The conservation and restoration of wild bees. *Ann. N. Y. Acad. Sci.* 1195, 169–197.
- Zanni, G., Opitz, J.M., 2013. Annals of morphology. Atavisms: phylogenetic Lazarus? *Am. J. Med. Genet.* 161, 2822–2835.