

# **FORMICA**

# Red Clover (2n)

Trifolium pratense L.

#### Things to know

The cultivar FORMICA has been systematically selected for a reduced content of formonenetin. This substance is supposed to be responsible for the oestrogenic effect of red clover on ruminants, which can affect their fertility. With the utilisation of FORMICA the risk of undesirable influences on animals can nearly be excluded, even with a high forage proportion of red clover. The remaining agronomic characteristics of FORMICA conform to the persistent "Mattenklee" type. However, the productivity of the variety can decrease in the second harvest year, especially with the appearance of anthracnose (Colletotrichum trifolii).

#### **Descent**

## Base material

Selection of landraces (collections 1970-1972). Selected on a reduced content of formonenetin.

# M0 seed

Row trials 1988 and 1990 with seed harvest on spaced plants progenies.

# Literature

Frick R., Jeangros B., Demenga M., Suter D., Hirschi H.-U., 2008. Essais de variétés de trèfle violet. Revue suisse Agric. 40(6), 245-248

Boller B., Tanner P., Schubiger F.-X., 2004. Merula und Pavo: neue, ausdauernde Mattenkleesorten. Agrarforschung 11(5), 162-167

# **National listing**

Situation in Switzerland

On the Swiss List of Recommended Varieties of Forage Plants since 1993

Situation abroad

FR (Representative: Jouffray-Drillaud, Cissé)

## **Agronomic caracteristics**

Results of the official Swiss variety trials 2005-2007

(Frick et al. 2008) ("Mattenklee" 2n) **FORMICA** Mean Yield 4.9 4.1 Juvenile growth 3.0 3.1 General impression 3.8 3.4 Competing ability 4.3 4.3 Persistence 5.1 4.3 3.4 Resistance to winter conditions 24 2.5 Resistance to Stemphylium Leafspots Resistance to black blotch and powdery mildew 1.7 Resistance to Anthracnose 2.3 1.8 Index (weighted average of all notes) 3.7 3.3

Scoring scale 1 = very good; 5 = medium; 9 = very poor Yield Mean of 5 experimental sites over 2 years Mean value of standard varieties

# **Description according to UPOV gidelines**

DUS test conducted at Scharnhorst, BSA (DE), 1995-1997

| UPOV No | Characteristics                             | State of expression        | Note |
|---------|---|----------------------------|------|
| 2       | Ploidy                                      |                            | 2    |
| 5       | Plant: natural height in the year of sowing | tall                       | 7    |
| 6       | Leaf: color in the year of sowing           | medium green to dark green | 6    |
| 9       | Plant: natural height in spring             | medium to tall             | 6    |
| 10      | Leaf: Color in spring                       | medium green               | 5    |
| 11      | Time of flowering                           | very early to early        | 2    |
| 12      | Stem: length                                | short                      | 3    |
| 14      | Stem: number of internodes                  | low to medium              | 4    |
| 16      | Leaf: shape of medial leaflet               | ovate                      | 2    |
|         |   |                            |      |

Version: 06.06.2012

Publisher: Agroscope Reckenholz-Tänikon Research Station ART,

Reckenholzstrasse 191, 8046 Zürich

In Collaboration with: Delley Seeds and Plants Ltd (DSP), 1567 Delley Editorial Team: Beat Boller, Peter Tanner, Lucienne Adam, ART

Copyright: © 2012, Art





Schweizerische Eidgenossenschaft Confédération suisse Confederazione Svizzera Confederaziun svizra

Swiss Confederation

Federal Department of Economic Affairs FDEA Agroscope Reckenholz-Tänikon Research Station ART