

OPEN ACCESS

EDITED BY

Luca Muzzioli,
Sapienza University of Rome, Italy

REVIEWED BY

Francesco Arfelli,
University of Bologna, Italy

*CORRESPONDENCE

Flaminia Ortenzi
✉ flamyortenzi@gmail.com

¹These authors share first authorship

RECEIVED 24 December 2025

REVISED 05 February 2026

ACCEPTED 06 February 2026

PUBLISHED 18 February 2026

CITATION

McAuliffe GA, Ortenzi F, van der Pols JC, Nemecek T, Colston J and Beal T (2026) Nutritional life cycle assessment for healthy and sustainable food systems: evidence and policy insights from Africa and Asia.

Front. Nutr. 13:1774865.

doi: 10.3389/fnut.2026.1774865

COPYRIGHT

© 2026 McAuliffe, Ortenzi, van der Pols, Nemecek, Colston and Beal. This is an open-access article distributed under the terms of the [Creative Commons Attribution License \(CC BY\)](#). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.

Nutritional life cycle assessment for healthy and sustainable food systems: evidence and policy insights from Africa and Asia

Graham A. McAuliffe^{1†}, Flaminia Ortenzi^{2*†},
Jolieke C. van der Pols³, Thomas Nemecek⁴, Jessica Colston⁵
and Ty Beal⁶

¹Harper Food Innovation, Harper Adams University, Newport, United Kingdom, ²Knowledge Leadership, Global Alliance for Improved Nutrition (GAIN), Geneva, Switzerland, ³School of Exercise and Nutrition Sciences and the Centre for Agriculture and the Bioeconomy, Queensland University of Technology (QUT), Brisbane, QLD, Australia, ⁴Life Cycle Assessment Research Group, Agroscope, Zurich, Switzerland, ⁵Programme Services Team, Global Alliance for Improved Nutrition (GAIN), London, United Kingdom, ⁶Knowledge Leadership, Global Alliance for Improved Nutrition (GAIN), Washington, DC, United States

Integrating nutritional value into Life Cycle Assessment (LCA) is essential for developing food system policies and interventions that simultaneously address environmental sustainability and human health. This Perspective explores recent conceptual and empirical evolutions in nutritional LCA (nLCA), drawing on expert talks, interdisciplinary stakeholder deliberations, and case studies presented at the 23rd International Union of Nutritional Sciences – International Congress of Nutrition, held in Paris in August 2025. We discuss methodological frameworks for incorporating nutritional quality into environmental footprint modelling, focusing on the selection of functional units and application of holistic nutrient profiling systems, such as the Nutritional Value Score. Case studies from Africa and Asia demonstrate the utility of nLCA to identify highly nutritious, lower-impact foods that mass- or energy-based denominators often overlook under attributional LCA. We argue that while plant-source foods frequently exhibit lower footprints, certain animal-source foods (such as small fish, dairy, eggs, and organ meats) can also be competitive when evaluated per unit of nutritional value. Finally, we highlight persistent challenges, including regional data gaps, lack of harmonisation in nutritional functional units, scope limitations, and risks of overinterpreting small differences in impact scores. While methodological refinement is still required, we conclude that nLCA offers a promising route for aligning agricultural, health, and environmental objectives, facilitating the development of more coherent food systems policies and programmes.

KEYWORDS

environmental impact, food policy, low- and middle-income countries, nLCA, nutrient profiling, nutritional life cycle assessment, sustainable food systems

1 Introduction

The urgent need to align environmental stewardship with global health priorities has recently placed food systems transformation at the centre of international policy debates (1–3). Life Cycle Assessment (LCA) often serves as a prominent methodology for quantifying the environmental burdens of goods or services. However, when applied to food systems, conventional mass- or energy-based LCA may yield incomplete, potentially misleading insights for decision-makers because it fails to account for the ultimate purpose of food: delivering adequate nutrition (4). We argue that for LCA to serve as a meaningful tool for food systems policy and interventions, it must be adapted to incorporate nutritional value – a framework increasingly recognised as nutritional LCA (nLCA) (4).

This perspective draws on evidence presented at a dedicated symposium on the evolution of nLCA within food systems held during the 23rd International Union of Nutritional Sciences – International Congress of Nutrition (IUNS-ICN) in Paris (August 2025; <https://www.icn2025.org/>). The session convened transdisciplinary and cross-sectoral experts to address *both* methodological advances *and* persistent challenges in applying nLCA to food and dietary assessments, with a particular focus on low- and middle-income country (LMIC) settings.

The symposium underscored that the move towards nLCA is not merely a technical adjustment but a necessary conceptual shift that allows for the prioritisation of supply chains, foods, and dietary patterns in ways that balance environmental goals with improved health and nutrition outcomes. The invited speakers covered the theoretical foundations of nLCA (4); the critical role of functional unit selection in determining the results (5–11); the use of holistic nutrient profiling systems such as the Nutritional Value Score (NVS) in environmental impact modelling (12); the application of combined *enviro-nutritional* footprint scores to local policy and programmatic decision-making (13); and current evidence gaps, limitations, and recommendations for future research (4, 14).

While significant methodological issues and data availability constraints remain, empirical case studies from diverse contexts – spanning LMIC (including, for instance, Indonesia, Kenya, and Rwanda) and high-income country (HIC; such as New Zealand, the United Kingdom, and Switzerland) settings – illustrate that nLCA can effectively reveal key trade-offs and synergies between nourishing a growing global population and minimising food-related environmental burdens (8, 13, 15, 16). By shifting the analytical lens from mass or energy to nutritional quality, nLCA offers a powerful framework for optimising food systems toward enhancing both human and planetary health.

2 The case for nutritional functional units

Designing healthier and more sustainable food systems requires effectively integrating environmental impacts with nutritional needs (1–3). Thus, the symposium opened with a strong rationale for nLCA as a data-driven tool to reconcile these often-conflicting objectives. Of note, large variability in nutritional and environmental priorities across geographic regions necessitates the selection of appropriate functional units that reflect the intended purpose of an

assessment – whether mass- or calorie-based, nutrient-specific (e.g., protein, calcium, vitamin A), or composite nutritional indices (e.g., NVS and, more commonly, the Nutrient Rich Food Index) (4–11). Crucially, the chosen metric must be relevant to the specific study context, taking into account the target population's nutritional requirements and the range and types of foods being compared. To this end, tailored weighting systems can be applied to composite indices to better reflect local needs (8, 17).

The NVS, a novel nutrient profiling system specifically designed for sustainability assessments, incorporates both nutrients of global health priority and dietary factors predictive of noncommunicable disease (NCD) risk, and adjusts for protein digestibility and bioavailability of iron and zinc – making it suitable for use in HIC and LMIC settings alike (12). When used as a functional unit, the NVS enables more nutritionally relevant comparisons of foods' environmental footprints than traditional mass-, energy-, or single-nutrient-based denominators (e.g., 1 kg, 1,000 Calories, or 100 g of protein). By holistically capturing foods' nutritional value, the NVS positions itself as a promising index for quantifying foods' environmental impacts relative to their contribution to global dietary adequacy and quality. For example, white rice and fatty fish have widely different nutrient profiles and play distinct roles in diets (i.e., source of energy from carbohydrates vs. source of protein, omega-3 s, and micronutrients); therefore, directly comparing their environmental footprints on a per-kilogram basis is not nutritionally robust. Even when comparing food items within the *same* category (e.g., two vegetables like spinach and carrots), mass- or energy-based metrics can yield misleading conclusions regarding their *enviro-nutritional* efficiency. For instance, a significantly smaller quantity of spinach (~250 g) achieves an equivalent NVS to a larger portion of carrots (~400 g), highlighting the difference in nutrient density and NCD-protective effects between these foods (Figure 1) (13). Further, taken from the perspective of LCA best practice (ISO, 2006), the *function* of distinct food items (within or across categories) at the point of consumption cannot be captured by mass, and only partially by energy, without accounting for broader nutritional considerations in parallel (18).

Empirical applications of nLCA in Indonesia, Kenya, and Rwanda demonstrated how locally adapted data and context-specific impact category weights can identify 'best-bet' foods for delivering high nutritional value at relatively low environmental costs. Our analyses revealed that while plant-source foods often perform well, certain minimally processed, nutrient-dense foods of animal origin – including organ meats, fish and seafood, eggs, and some dairy products – can be competitive when assessed per unit of nutritional value, particularly if consumed in line with national dietary recommendations (13). These findings challenge simplistic dichotomies and point to the need for nuanced, context-sensitive guidance regarding consumption of plant and animal products within healthy and sustainable diets (4–11).

3 Current challenges and implications for decision-making

From a methodological perspective, we highlight significant limitations surrounding functional unit selection, assessment scope, approach standardisation, transparency, replicability, results interpretation, and uncertainty communication (4–11).




FIGURE 1

Amount (grams) of common Indonesian foods needed to achieve equivalent Nutritional Value Scores – in this case, an NVS of 100. The NVS rates foods on a scale of 1 to 100, where 1 represents the least nutritious food and 100 the most nutritious one (12). While this ranking facilitates both within- (e.g., different vegetables) and cross-category (e.g., vegetables vs. starchy staples) comparisons of foods' nutritional value, it does not imply direct interchangeability of items in meals or diets. Foods play distinct roles in a diet, and all recommended food groups consumed in adequate proportions (i.e., as per dietary guidelines) should be part of healthy and sustainable eating patterns. Reproduced from Ortenzi et al (13), licensed under CC BY-NC-SA 4.0 IGO.

Unlike other recent adaptations of, and advances in LCA, such as prospective and spatial LCA (19–22), nutritional LCA can introduce health-based narratives which feed into, either directly or indirectly, recommendations for individual- and/or population-level dietary changes (23, 24). The proposed consumption shifts may have significant implications for nutrition and health outcomes. Therefore, our symposium stressed the importance of scientific and clinical due diligence, as well as local health professional engagement, and called for caution against overinterpreting small differences in *enviro-nutritional* footprint scores given the inherent uncertainties in (n)LCA modeling (25).

The utility of nLCA-derived insights extends across policy, programmatic, and industrial spheres, from informing sustainable dietary guidelines and optimised public procurement strategies, to competitive product benchmarking, consumer-demand generation initiatives, and the identification of environmental 'hotspots' throughout supply chains (Figure 2) (13). For example, recent work led by the Global Alliance for Improved Nutrition in Indonesia illustrated how nLCA findings have already been leveraged to influence (sub)national food systems policy and action planning, and to incentivise innovation in local food production (13, 26, 27). When it comes to informing public and corporate

decision-making, clear results communication is essential: while aggregated *enviro-nutritional* impact scores can aid interpretation by diverse (technical and non-) stakeholders, they risk obscuring the relative importance of and potential trade-offs among individual environmental categories. This concern is of notable relevance for spatially-dependent impact categories (e.g., water and soil pollution potentials, including eutrophication and acidification) (4, 13).

In addition to these conceptual and interpretative issues, there are persistent, wide-ranging data gaps, particularly regarding the availability and geographic representativeness of both nutritional and environmental datasets, which are unevenly distributed across regions, value chains, and production systems (4–11).

Current evidence emphasises that transdisciplinary and multisectoral collaboration is essential to improve data quality and spatial granularity, as well as to harmonise methods and ensure appropriate use of results, toward enhancing local relevance of analytical outputs, enabling cross-study comparisons, and facilitating coherent policy and intervention design in food systems (4–11).

4 Discussion and future directions in *enviro-nutritional* modelling

The open panel discussion that followed the session's structured presentations reflected the diverse composition of the audience, with contributions from public health professionals, nutrition experts, environmental scientists, policymakers, programme managers, and private sector representatives. Questions centred on the feasibility of applying complex nutrient profiling systems and nLCA models in data-scarce settings; the poorly captured role of cultural acceptability and socio-economic impacts in defining sustainable diets; and the potential for nLCA to guide reformulation and innovation in the food industry.

Reflecting the symposium's consensus, we suggest that advancing nLCA requires progress in three key areas to unlock its full potential as an evidence-based tool to inform high-stakes food system decision-making. In the *nutritional domain*, methodological refinement must prioritise (i) developing practically viable yet comprehensive nutrient indices that are fit-for-purpose and contextually relevant, and (ii) improving the quality and regional coverage of dietary intake and food composition datasets. Simultaneously, research in the *environmental domain* must also address data infrastructure gaps, as well as expand assessment scopes – e.g., to include a broader range of value chains, production systems, and impact categories – and parameter-level uncertainty measurements. Building on these disciplinary foundations, future *integrated assessments* should focus on (i) harmonising functional units and analytical approaches, and (ii) incorporating other sustainability dimensions (i.e., socio-economic and cultural) within the same LCA framework, to generate consistent, transparent, and holistic metrics (4–11, 17).

In alignment with the broader literature, all invited experts and session participants emphasised that the current nLCA evidence base is heavily skewed toward HICs. Closing the 'data equity' gap – particularly across Sub-Saharan Africa and parts of Asia – is essential for ensuring that nLCA results reflect local agroecological conditions, production practices, and food consumption patterns. Failure to improve geographic granularity may lead to 'one-size-fits-all' policy




FIGURE 2

Contribution of different lifecycle stages to climate change impacts (expressed as kg CO₂ eq/100 NVS) for a selection of foods commonly consumed in Indonesia. The NVS (specifically, an NVS of 100) was used as the nutritional functional unit for this analysis (12). Reproduced from Ortenzi et al (13), licensed under CC BY-NC-SA 4.0 IGO.

decisions in LMICs based on models derived from HIC archetypes (4–11).

Moreover, future nLCA efforts should evolve beyond attributional, climate-centric assessments to embrace prospective and geo-spatial modelling (19, 20). This transition will allow researchers to estimate the nutrient provision capacity of alternative farming systems relative to the arable land required, offering a more sophisticated understanding of land-use efficiency (21, 22).

Further developments are critically needed to consider broader sustainability aspects – such as biodiversity loss, animal welfare, and the sociocultural and livelihoods implications of supply chains – and to explore meal- and whole diet-level (in addition to product-level) nLCA models, as well as system-wide evaluations where food production and consumption perspectives are integrated. For instance, applying nLCA methodology to pre-defined dietary scenarios (e.g., national average consumption patterns, dietary guidelines, school canteen menus) may offer more comprehensive, policy-relevant insights than single-product assessments, as it inherently accounts for nutritional complementary of food items while measuring environmental and social impacts (4, 28–30). Nevertheless, granular comparisons both within and across food groups remain essential for benchmarking specific commodities and guiding targeted food-based interventions.

The first-time inclusion of a dedicated nLCA symposium at such a prominent nutrition science conference signals significant progress toward mainstreaming environmental footprint evaluations in global food systems research and policy agendas. However, to prevent the

fragmentation and incomparability of findings from different studies, there is an urgent need to harmonise nutritional functional units and establish transparent reporting standards. This is especially important given the recently introduced mandatory reporting policies in many countries and regions, such as the European Union's Corporate Sustainability Reporting Directive and the Australian Sustainability Reporting Standards (31, 32).

5 Conclusion

By combining environmental burden and nutritional value quantification within the same analytical frame, nLCA offers a route to more coherent and holistic food systems policies and interventions. The evidence synthesised in this Perspective illustrates that nLCA represents a major methodological advancement from mass- or energy-based assessments, which often penalise nutrient-rich foods with great potential to positively contribute to global dietary adequacy and quality. By shifting to nutrition-based denominators, we can better characterise the synergistic effects of 'best-bet' foods that support both human and planetary health.

While methodological limitations remain significant, the momentum established at the IUNS-ICN 2025 session demonstrates a clear appetite for interdisciplinary alignment. The challenge for the coming decade is not merely technical, but also ethical and political: we must leverage *enviro-nutritional* modelling in ways that respect local

sociocultural realities and promote economic prosperity while contributing to global sustainability and health targets.

Ultimately, embedding nLCA within the food and nutrition science field is no longer a 'niche' research interest; it is a requirement for designing future-proof food systems. For researchers, policymakers, and industry actors alike, nLCA can offer the cross-sectoral language necessary to reconcile the often-conflicting goals of agricultural production, environmental protection, and public health nutrition.

Data availability statement

Publicly available datasets were analyzed in this study. This data can be found here: all data referred to in this perspective can be found in previous publications (see citations) and in the following repository: McAuliffe, G, Ortenzi, F, McLaren, S, Ponsioen, T, Beal, T. Underlying Nutritional Life Cycle Assessment (nLCA) inventory data applicable to an Indonesian case study of the Nutritional Value Score (NVS). Mendeley Data. 2025; V1. Available at: <https://data.mendeley.com/datasets/2fzwjty3jc/1>.

Author contributions

GM: Conceptualization, Data curation, Formal analysis, Investigation, Methodology, Software, Writing – original draft, Writing – review & editing. FO: Conceptualization, Data curation, Formal analysis, Investigation, Methodology, Project administration, Writing – original draft, Writing – review & editing. JP: Writing – review & editing. TN: Writing – review & editing. JC: Project administration, Writing – review & editing. TB: Conceptualization, Formal analysis, Investigation, Methodology, Software, Visualization, Writing – review & editing.

Funding

The author(s) declared that financial support was received for this work and/or its publication. The nLCA symposium at IUNS-ICN 2025 was sponsored by the Global Alliance for Improved Nutrition (GAIN) via the Nourishing Food Pathways programme, jointly funded by the German Federal Ministry for Economic Cooperation and Development; the Ministry of Foreign Affairs of the Netherlands; the

References

- Food and Agriculture Organization of the United Nations (FAO). Sustainable food systems. concept and framework. Rome, Italy: Food and Agriculture Organization of the United Nations (FAO); 2018. Available online at: <https://openknowledge.fao.org/server/api/core/bitstreams/b620989c-407b-4caf-a152-f790f55fec71/contents>
- High Level Panel of Experts on Food Security and Nutrition (HLPE). Nutrition and food systems. a report by the high level panel of experts on food security and nutrition of the committee on world food security. Rome, Italy: High Level Panel of Experts on Food Security and Nutrition (HLPE); 2017. Available online at: <https://openknowledge.fao.org/server/api/core/bitstreams/4ac1286e-eef3-4f1d-b5bd-d92f5d1ce738/content>
- Rockström J, Thilsted SH, Willett WC, Gordon LJ, Herrero M, Hicks CC, et al. The EAT-lancet commission on healthy, sustainable, and just food systems. *Lancet.* (2025) 406:1625–700. doi: 10.1016/S0140-6736(25)01201-2
- McLaren S, Berardy A, Henderson A, Holden N, Huppertz T, Jolliet O, et al. Integration of environment and nutrition in life cycle assessment of food items: Opportunities and challenges. Food and Agriculture Organization of the United Nations (FAO); 2021 (Accessed 2025 Jun 13). Available online at: <https://openknowledge.fao.org/handle/20.500.14283/cb8054en>
- Green A, Nemecek T, Mathys A. A proposed framework to develop nutrient profiling algorithms for assessments of sustainable food: the metrics and their assumptions matter. *Int J Life Cycle Assess.* (2023) 28:1326–47. doi: 10.1007/s11367-023-02210-9
- Hallström E, Davis J, Woodhouse A, Sonesson U. Using dietary quality scores to assess sustainability of food products and human diets: a systematic review. *Ecol Indic.* (2018) 93:219–30. doi: 10.1016/j.ecolind.2018.04.071
- Katz-Rosene R, Ortenzi F, McAuliffe GA, Beal T. Levelling foods for priority micronutrient value can provide more meaningful environmental footprint comparisons. *Commun Earth Environ.* (2023) 4:287. doi: 10.1038/s43247-023-00945-9

8. Majumdar S, McLaren SJ, van der Pols JC, Lister CE. An nLCA approach to support consumer meal decisions: a New Zealand case study of toppings on toast. *Front Sustain Food Syst.* (2024) 8:1363565. doi: 10.3389/fsufs.2024.1363565/full
9. McAuliffe GA, Takahashi T, Beal T, Huppertz T, Leroy F, Buttriss J, et al. Protein quality as a complementary functional unit in life cycle assessment (LCA). *Int J Life Cycle Assess.* (2023) 28:146–55. doi: 10.1007/s11367-022-02123-z
10. McAuliffe GA, Takahashi T, Lee MRF. Applications of nutritional functional units in commodity-level life cycle assessment (LCA) of Agri-food systems. *Int J Life Cycle Assess.* (2020) 25:208–21. doi: 10.1007/s11367-019-01679-7
11. Sonesson U, Davis J, Flysjö A, Gustavsson J, Withöft C. Protein quality as functional unit – a methodological framework for inclusion in life cycle assessment of food. *J Clean Prod.* (2017) 140:470–8. doi: 10.1016/j.jclepro.2016.06.115
12. Beal T, Ortenzi F. Nutritional value score rates foods based on global health priorities. Research Square; (2025). Available online at: <https://www.researchsquare.com/article/rs-3443927/v2>
13. Ortenzi F, Colston J, Beal T. *Nourishing people and planet: Enviro-nutritional insights into local foods for policy, Programmes, and industry.* Geneva, Switzerland: Global Alliance for Improved Nutrition (GAIN) (2025).
14. McAuliffe GA, Beal T, Lee MRF, van der Pols JC. Editorial: pushing the frontiers of nutritional life cycle assessment (nLCA) to identify globally equitable and sustainable Agri-food systems. *Front Sustain Food Syst.* (2024) 8:1471102. doi: 10.3389/fsufs.2024.1471102/full
15. Herrmann M, Mehner E, Egger L, Portmann R, Hammer L, Nemecek T. A comparative nutritional life cycle assessment of processed and unprocessed soy-based meat and milk alternatives including protein quality adjustment. *Front Sustain Food Syst.* (2024) 8:1413802. doi: 10.3389/fsufs.2024.1413802/full
16. McAuliffe GA, Takahashi T, Lee MRF. Framework for life cycle assessment of livestock production systems to account for the nutritional quality of final products. *Food Energy Secur.* (2018) 7:e00143. doi: 10.1002/fes3.143
17. Ridoutt B. An alternative nutrient rich food index (NRF-ai) incorporating prevalence of inadequate and excessive nutrient intake. *Foods* 2021;10. Available online at: <https://www.mdpi.com/2304-8158/10/12/3156>
18. ISO. ISO 14044:2006: environmental management — life cycle assessment — requirements and guidelines. 1st edn. (2006) (Accessed 2026 Feb 5). Available online at: <https://www.iso.org/standard/38498.html>
19. Arvidsson R, Svanström M, Sandén BA, Thonemann N, Steubing B, Cucurachi S. Terminology for future-oriented life cycle assessment: review and recommendations. *Int J Life Cycle Assess.* (2024) 29:607–13. doi: 10.1007/s11367-023-02265-8
20. Douziech M, Bauer C, Bystricky M, Diogo V, Jouannais P, Lansche J, et al. Prospective LCA to support the Agri-food sector's transition towards sustainability—89th LCA discussion forum conference report. *Int J Life Cycle Assess.* (2025) 30, 2794–2798. doi: 10.1007/s11367-025-02556-2
21. Lee MRF, Domingues JP, McAuliffe GA, Tichit M, Accatino F, Takahashi T. Nutrient provision capacity of alternative livestock farming systems per area of arable farmland required. *Sci Rep.* (2021) 11:14975. doi: 10.1038/s41598-021-93782-9
22. McAuliffe GA, Zhang Y, Collins AL. Assessing catchment scale water quality of Agri-food systems and the scope for reducing unintended consequences using spatial life cycle assessment (LCA). *J Environ Manag.* (2022) 318:115563. doi: 10.1016/j.jenvman.2022.115563
23. Stylianou KS, Fulgoni VL, Jolliet O. Small targeted dietary changes can yield substantial gains for human health and the environment. *Nat Food.* (2021) 2:616–27. doi: 10.1038/s43016-021-00343-4
24. Stylianou KS, Heller MC, Fulgoni VL, Ernstaff AS, Keoleian GA, Jolliet O. A life cycle assessment framework combining nutritional and environmental health impacts of diet: a case study on milk. *Int J Life Cycle Assess.* (2016) 21:734–46. doi: 10.1007/s11367-015-0961-0
25. Ortenzi F, McAuliffe GA, Leroy F, Nordhagen S, van Vliet S, del Prado A, et al. Can we estimate the impact of small targeted dietary changes on human health and environmental sustainability? *Environ Impact Assess Rev.* (2023) 102:107222. doi: 10.1016/j.eiar.2023.107222
26. de Pee S, Hardinsyah R, Jalal F, Kim BF, Semba RD, Deptford A, et al. Balancing a sustained pursuit of nutrition, health, affordability and climate goals: exploring the case of Indonesia. *Am J Clin Nutr.* (2021) 114:1686–97. doi: 10.1093/ajcn/nqab258
27. Sari E, Budiman I. Jack bean: A resilient legume to improve 'tempeh security' in Indonesia. (2025). Available online at: <https://www.gainhealth.org/blogs/jack-bean-resilient-legume-improve-tempeh-security-indonesia>
28. Campobasso V, Gallucci T, Crovella T, Vignali G, Paiano A, Lagioia G, et al. Life cycle assessment of food catering menus in a university canteen located in southern Italy. *Sci Total Environ.* (2024) 957:177482. doi: 10.1016/j.scitotenv.2024.177482
29. Arfelli F, Ciacci L, Cespi D, Vassura, Passarini F. The "SQUIID claim": a novel LCA-based indicator for food dishes. *J Clean Prod.* (2024) 434:140241. doi: 10.1016/j.jclepro.2023.140241
30. Yacoub Bach L, Jana BE, Adaeze Egwutu CF, Orndorff CJ, Alanakrih R, Okoro J, et al. A sustainability analysis of environmental impact, nutritional quality, and price among six popular diets. *Front Sustain Food Syst.* (2023) 7:1021906. doi: 10.3389/fsufs.2023.1021906/full
31. Australian Accounting Standards Board (AASB). Voluntary AASB S1 general requirements for disclosure of sustainability-related financial information and mandatory AASB S2 climate-related disclosures. 2024. Available online at: <https://aasb.gov.au/news/australian-sustainability-reporting-standards-aasb-s1-and-aasb-s2-are-now-available-on-the-aasb-digital-standards-portal/>
32. European Union. Directive (EU) 2022/2464 of the European Parliament and of the council of 14 December 2022 amending regulation (EU) no 537/2014, directive 2004/109/EC, directive 2006/43/EC and directive 2013/34/EU, as regards corporate sustainability reporting (corporate sustainability reporting directive). *Off J Eur Union*, 322, pp. 15–62; 2022. Available online at: https://eur-lex.europa.eu/legal-content/EN/TXT/?uri=CELE_X%3A32022L2464