

Fact Sheet

LACERTA

Perennial Ryegrass (4n)

Lolium perenne L.

Things to know

The tetraploid cultivar LACERTA has proved to be very successful, under many cultivation conditions in international comparison. Especially the high resistance to crown rust and the strong ability to grow again in summertime, distinguishes it from the other precocious cultivars. It recovers rapidly from winter damages and suppresses weed growth. LACERTA builds rather loose stands. It is easy to digest and due to its competitive quality, LACERTA contributes to a stable part of high quality grass in mixtures.

Descent

Base material

Selection of Arion material after colchicine treatment.

M0 seed

Row trials (LP8935/45) with seed harvest on 53 clone progenies from a polyccross 1987.

Literature

Suter D., Hirschi H.-U., Frick R., 2017. Englisches Raigras: neue Sorten für den Schweizer Kunstfutterbau empfohlen. Agrarforschung Schweiz 8(7), 292-299

Suter D., Hirschi H.-U., Frick R., Aebi P., 2012. Englisches Raigras: 62 Sorten mussten sich bewähren. Agrarforschung Schweiz 3(9), 414-421

National listing

Situation in Switzerland

On the Swiss List of Recommended Varieties of Forage Plants from 1996 to

Situation abroad

DE (Representative: Euro Grass, Lippstadt)

Agronomic caracteristics

Results of the official Swiss variety trials 2014-2016

| (Suter et al. 2017) (4n, früh) | LACERTA | Mean |
|---------------------------------------|---------|------|
| Yield | 4.9 | 3.9 |
| General impression | 3.4 | 2.8 |
| Juvenile growth | 1.9 | 1.7 |
| Competing ability | 5.0 | 5.0 |
| Persistence | 4.3 | 3.9 |
| Resistance to winter conditions | 4.9 | 4.4 |
| Resistance to leafspots an rust | 2.6 | 2.5 |
| Digestibility of the organic matter | 5.3 | 4.6 |
| Persistence at higher altitudes | 2.8 | 2.2 |
| Index (weighted average of all notes) | 3.7 | 3.2 |

Scoring scale 1 = very good; 5 = medium; 9 = very poor Yield Mean of 4 experimental sites over 2 years Mean Mean value of standard varieties

Description according to UPOV gidelines

DUS test conducted at Scharnhorst, BSA (DE), 1997-1999

| D 0 0 1001 0 | oridation at Contaminoral, Bort (BE), 1001 | 1000 | |
|--------------|---|-------------------------------|------|
| UPOV No | Characteristics | State of expression | Note |
| 1 | Ploidy | tetraploid | 4 |
| 5 | Leaf: color in the year of sowing | medium green to dark green | 6 |
| 8 | Time of inflorescence emergence (after vernalization) | very early to early | 2 |
| 10 | Flag leaf: length | medium to long | 6 |
| 11 | Flag leaf: width | broad | 7 |
| 12 | Plant: length of longest stem including inflorescence | long | 7 |

Version: 16.02.2018

Publisher: Agroscope, Reckenholzstrasse 191, 8046 Zürich In Collaboration with: Delley Seeds and Plants Ltd (DSP), 1567 Delley Editorial Team: Christoph Grieder and Peter Tanner, Agroscope

Copyright: © 2018, Agroscope



Schweizerische Eidgenossenschaft Confédération suisse Confederazione Svizzera Confederaziun svizra

Swiss Confederation

Federal Department of Economic Affairs, Education and Research EAER Agroscope